

HANKEL MULTIPLIERS OF LAPLACE TRANSFORM TYPE

J.J. BETANCOR, A.J. CASTRO, AND J. CURBELO

ABSTRACT. In this paper we prove that the Hankel multipliers of Laplace transform type on $(0, \infty)^n$ are of weak type $(1, 1)$. Also we analyze L^p -boundedness properties for the imaginary powers of Bessel operator on $(0, \infty)^n$.

1. INTRODUCTION

We consider the Hankel transformation on $(0, \infty)^n$ defined by

$$h_{\lambda_1, \dots, \lambda_n}(f)(x) = \int_{(0, \infty)^n} \prod_{j=1}^n (x_j y_j)^{-\lambda_j + 1/2} J_{\lambda_j - 1/2}(x_j y_j) f(y) \prod_{j=1}^n y_j^{2\lambda_j} dy, \quad x \in (0, \infty)^n,$$

where $\lambda_j > -1/2$, $j = 1, \dots, n$. Here J_ν denotes the Bessel function of the first kind and order ν . $h_{\lambda_1, \dots, \lambda_n}$ maps $L^1((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$ into $L^\infty((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$ ($= L^\infty((0, \infty)^n, dx)$) and it can be extended as an isometry to $L^2((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$ being $h_{\lambda_1, \dots, \lambda_n}^{-1} = h_{\lambda_1, \dots, \lambda_n}$ on $L^2((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$ (see [3]).

If $m \in L^\infty((0, \infty)^n, dx)$ the multiplier operator $T_{\lambda_1, \dots, \lambda_n}^m$ associated with the Hankel transformation $h_{\lambda_1, \dots, \lambda_n}$ is defined by

$$T_{\lambda_1, \dots, \lambda_n}^m(f) = h_{\lambda_1, \dots, \lambda_n}(m h_{\lambda_1, \dots, \lambda_n}(f)), \quad f \in L^2((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx).$$

It is clear that $T_{\lambda_1, \dots, \lambda_n}^m$ is bounded from $L^2((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$ into itself. L^p -boundedness properties for Hankel multipliers have been studied in [1], [2], [6], [7], [8], [9], [10], [16] and [21], amongst others.

In this paper, following the ideas presented in [5] and in [17] we study Hankel multiplier operator when m is of Laplace transform type. For every $\lambda > -1/2$ we denote by $\Delta_{\lambda, x}$ the Bessel operator

$$\Delta_{\lambda, x} = -x^{-2\lambda} \frac{d}{dx} \left(x^{2\lambda} \frac{d}{dx} \right) = - \left(\frac{d^2}{dx^2} + \frac{2\lambda}{x} \frac{d}{dx} \right),$$

and if $\lambda_j > -1/2$, $j = 1, \dots, n$, we write

$$\Delta_{\lambda_1, \dots, \lambda_n} = \sum_{j=1}^n \Delta_{\lambda_j, x_j}.$$

Since

$$\Delta_{\lambda, x} \left[(xy)^{-\lambda+1/2} J_{\lambda-1/2}(xy) \right] = y^2 (xy)^{-\lambda+1/2} J_{\lambda-1/2}(xy), \quad x, y \in (0, \infty)$$

(see [11, (5.3.7), p. 103]), it follows that, for each $x = (x_1, \dots, x_n)$, $y = (y_1, \dots, y_n) \in (0, \infty)^n$,

$$(1) \quad \Delta_{\lambda_1, \dots, \lambda_n} \left[\prod_{j=1}^n (x_j y_j)^{-\lambda_j + \frac{1}{2}} J_{\lambda_j - \frac{1}{2}}(x_j y_j) \right] = \left(\sum_{j=1}^n y_j^2 \right) \prod_{j=1}^n (x_j y_j)^{-\lambda_j + \frac{1}{2}} J_{\lambda_j - \frac{1}{2}}(x_j y_j).$$

This paper is partially supported by MTM2007/65609.

According to (1) we say that m is of Laplace transform type when

$$(2) \quad m(y) = \sum_{j=1}^n y_j^2 \int_0^\infty e^{-t \sum_{j=1}^n y_j^2} \phi(t) dt, \quad y \in (0, \infty)^n,$$

for a certain $\phi \in L^\infty(0, \infty)$ ([17, p. 121]).

Our main result is the following.

Theorem 1.1. *Let $\lambda_j > -1/2$, $j = 1, \dots, n$. Assume that m is of Laplace transform type. Then, the Hankel multiplier $T_{\lambda_1, \dots, \lambda_n}^m$ is bounded from $L^p((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$ into itself, for every $1 < p < \infty$, and from $L^1((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$ into $L^{1,\infty}((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$.*

This theorem will be proved in Section 2. It can be seen as an extension to higher dimension of [1, Theorem 1.2]. However, in order to prove Theorem 1.1 we use a different procedure than the one employed in the proof of [1, Theorem 1.2]. In [1] Calderón-Zygmund theory for singular integral operators is applied. Here, we represent the multiplier operator $T_{\lambda_1, \dots, \lambda_n}^m$ as a principal value integral operator when it acts on the space $C_c^\infty((0, \infty)^n)$ of the C^∞ functions with compact support in $(0, \infty)^n$ (see Proposition 2.1). Then, after proving L^p -boundedness properties for the maximal operator associated with the principal value integral operator (see Proposition 2.2), we extend the Hankel multiplier $T_{\lambda_1, \dots, \lambda_n}^m$ to $L^p((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$, $1 \leq p < \infty$, as a principal value integral operator that is bounded from $L^p((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$ into itself, when $1 < p < \infty$, and from $L^1((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$ into $L^{1,\infty}((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$ (see Proposition 2.3). To establish these properties we split the region $(0, \infty)^n \times (0, \infty)^n$ in two parts. The set

$$\Omega = \{(x, y) \in (0, \infty)^n \times (0, \infty)^n : x_j/2 < y_j < 2x_j, j = 1, \dots, n\}$$

is called the local region. In Ω the kernel $K_{\lambda_1, \dots, \lambda_n}^\phi$ that defines the Hankel multiplier $T_{\lambda_1, \dots, \lambda_n}^m$ differs from $\prod_{j=1}^n (x_j y_j)^{-\lambda_j} K^m$, where K^m is the kernel associated with the Fourier multipliers T^m defined by

$$T^m f = (m\hat{f})^\vee, \quad f \in L^2(\mathbb{R}, dx),$$

by a kernel defining a bounded operator in $L^p((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$, for every $1 \leq p < \infty$. Here as usual, by $\hat{f}$ we denote the Fourier transform of f and by $\check{g}$ the inverse Fourier transform of g .

On $(0, \infty)^n \times (0, \infty)^n \setminus \Omega$, called global region, the kernel $|K_{\lambda_1, \dots, \lambda_n}^\phi|$ defines a positive bounded operator from $L^p((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$ into itself, when $1 < p < \infty$, and from $L^1((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$ into $L^{1,\infty}((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$.

The semigroup $\{W_t^{\lambda_1, \dots, \lambda_n}\}_{t \geq 0}$ generated by $-\Delta_{\lambda_1, \dots, \lambda_n}$ is defined as follows

$$W_t^{\lambda_1, \dots, \lambda_n}(f)(x) = \int_{(0, \infty)^n} \prod_{j=1}^n W_t^{\lambda_j}(x_j, y_j) f(y) \prod_{j=1}^n y_j^{2\lambda_j} dy, \quad x = (x_1, \dots, x_n) \in (0, \infty)^n,$$

where

$$(3) \quad W_t^\lambda(x, y) = \frac{(xy)^{-\lambda+1/2}}{2t} I_{\lambda-1/2}\left(\frac{xy}{2t}\right) e^{-(x^2+y^2)/4t}, \quad t, x, y \in (0, \infty), \lambda > -1/2,$$

and I_ν denotes the modified Bessel function of the first kind and order ν ([22, p. 395]). This semigroup $\{W_t^{\lambda_1, \dots, \lambda_n}\}_{t \geq 0}$ is a symmetric diffusion semigroup in the sense of Stein ([17, p. 65]). Moreover, the Hankel multiplier $T_{\lambda_1, \dots, \lambda_n}^m$ is actually an spectral multiplier associated with $\Delta_{\lambda_1, \dots, \lambda_n}$.

Then, by [17, Corollary 3, p. 121] $T_{\lambda_1, \dots, \lambda_n}^m$ is bounded from $L^p((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$ into itself, for every $1 < p < \infty$. In Theorem 1.1 we prove as a new result that $T_{\lambda_1, \dots, \lambda_n}^m$ defines a bounded operator from $L^1((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$ into $L^{1,\infty}((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$. Moreover, in Proposition 2.3 we establish a representation of the operator $T_{\lambda_1, \dots, \lambda_n}^m$ as a principal value integral operator in $L^p((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$, $1 \leq p < \infty$.

As an application of Theorem 1.1 we can show L^p -boundedness properties for the imaginary powers of the Bessel operator $\Delta_{\lambda_1, \dots, \lambda_n}$. We define, for every $\beta \in \mathbb{R}$, the function

$$\phi_\beta(t) = \frac{t^{-i\beta}}{\Gamma(1-i\beta)}, \quad t \in (0, \infty).$$

We have that

$$m_\beta(y) = \sum_{j=1}^n y_j^2 \int_0^\infty e^{-t \sum_{j=1}^n y_j^2} \phi_\beta(t) dt = \left(\sum_{j=1}^n y_j^2 \right)^{i\beta}, \quad y = (y_1, \dots, y_n) \in (0, \infty)^n \text{ and } \beta \in \mathbb{R}.$$

For every $\beta \in \mathbb{R}$, the $i\beta$ -power $\Delta_{\lambda_1, \dots, \lambda_n}^{i\beta}$ of $\Delta_{\lambda_1, \dots, \lambda_n}$ is defined by

$$\Delta_{\lambda_1, \dots, \lambda_n}^{i\beta} = T_{\lambda_1, \dots, \lambda_n}^{m_\beta}.$$

From Theorem 1.1 we deduce the following.

Corollary 1.1. *Let $\beta \in \mathbb{R}$. Then, the operator $\Delta_{\lambda_1, \dots, \lambda_n}^{i\beta}$ is bounded from $L^p((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$ into itself, for every $1 < p < \infty$, and from $L^1((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$ into $L^{1,\infty}((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$.*

The next properties of the modified Bessel function I_ν , $\nu > -1$, that can be found in [11, Chapter 5], are very useful in the sequel:

$$(4) \quad z^{-\nu} I_\nu(z) \sim \frac{1}{2^\nu \Gamma(\nu+1)}, \quad \text{as } z \rightarrow 0^+;$$

$$(5) \quad I_\nu(z) = \frac{e^z}{\sqrt{2\pi z}} \left(\sum_{k=0}^n (-1)^k [\nu, k] (2z)^{-k} + \mathcal{O}\left(\frac{1}{z^{n+1}}\right) \right), \quad n \in \mathbb{N}, \quad z \in (0, \infty),$$

where $[\nu, 0] = 1$ and

$$[\nu, k] = \frac{(4\nu^2 - 1)(4\nu^2 - 3^2) \dots (4\nu^2 - (2k-1)^2)}{2^{2k} \Gamma(k+1)}, \quad k = 1, 2, \dots;$$

and

$$(6) \quad \frac{d}{dz} (z^{-\nu} I_\nu(z)) = z^{-\nu} I_{\nu+1}(z), \quad z \in (0, \infty).$$

Also, we will use the following properties of the Bessel function J_ν , $\nu > -1$, (see [11, pages 110 and 123]):

$$(7) \quad z^{-\nu} J_\nu(z) \sim \frac{1}{2^\nu \Gamma(\nu+1)}, \quad \text{as } z \rightarrow 0^+;$$

$$(8) \quad J_\nu(z) = \sqrt{\frac{2}{\pi z}} \left(\cos\left(z - \frac{\nu\pi}{2} - \frac{\pi}{4}\right) + \mathcal{O}\left(\frac{1}{z}\right) \right), \quad z \in (0, \infty);$$

and

$$(9) \quad \frac{d}{dz} (z^{-\nu} J_\nu(z)) = -z^{-\nu} J_{\nu+1}(z), \quad z \in (0, \infty).$$

The heat kernel associated with the Bessel operator $-\Delta_{\lambda, x}$ can be written in terms of Bessel functions by ([22, p. 395])

$$(10) \quad W_t^\lambda(x, y) = \int_0^\infty e^{-tz^2} (zx)^{-\lambda+1/2} J_{\lambda-1/2}(zx) (zy)^{-\lambda+1/2} J_{\lambda-1/2}(zy) z^{2\lambda} dz, \quad t, x, y \in (0, \infty).$$

The n -dimensional Hankel transform on function and distribution spaces has been studied by Molina and Trione ([12]). A formula connecting Hankel transform and Bessel operators that we will use is the following

$$(11) \quad h_{\lambda_1, \dots, \lambda_n}(\Delta_{\lambda_1, \dots, \lambda_n} f)(x) = |x|^2 h_{\lambda_1, \dots, \lambda_n}(f)(x), \quad x \in (0, \infty)^n \text{ and } f \in C_c^\infty((0, \infty)^n).$$

Also the Plancherel type equality

$$(12) \quad \int_{(0, \infty)^n} h_{\lambda_1, \dots, \lambda_n}(f)(x) h_{\lambda_1, \dots, \lambda_n}(g)(x) \prod_{j=1}^n x_j^{2\lambda_j} dx = \int_{(0, \infty)^n} f(x) g(x) \prod_{j=1}^n x_j^{2\lambda_j} dx,$$

holds for every $f, g \in L^2((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$ (see [3, Lemma 2.7]).

Throughout this paper we will always denote by C a suitable positive constant that can change from one line to the other one. Also, we will use repeatedly without saying it that, for every $k \in \mathbb{N}$, $\sup_{z>0} z^k e^{-z} < \infty$.

2. PROOF OF THEOREM 1.1

In order to prove Theorem 1.1 we need to establish a pointwise integral representation for the multiplier operator $T_{\lambda_1, \dots, \lambda_n}^m$ as a principal value integral operator (see Proposition 2.1 below). In the sequel we assume that

$$m(x) = \sum_{j=1}^n x_j^2 \int_0^\infty e^{-t \sum_{j=1}^n x_j^2} \phi(t) dt, \quad x \in (0, \infty)^n,$$

where $\phi \in L^\infty((0, \infty)^n)$.

Firstly we prove the following result.

Lemma 2.1. *Let $\lambda_j > -1/2$, $j = 1, \dots, n$. Assume that $f \in C_c^\infty((0, \infty)^n)$. Then,*

$$T_{\lambda_1, \dots, \lambda_n}^m(f)(x) = \int_0^\infty \phi(t) h_{\lambda_1, \dots, \lambda_n} \left(\sum_{j=1}^n y_j^2 e^{-t \sum_{j=1}^n y_j^2} h_{\lambda_1, \dots, \lambda_n}(f)(y) \right) (x) dt, \quad a.e. \ x \in (0, \infty)^n.$$

Proof. Let $g \in C_c^\infty((0, \infty)^n)$. By using Plancherel equality (12) we get

$$\begin{aligned} & \int_{(0, \infty)^n} T_{\lambda_1, \dots, \lambda_n}^m(f)(x) g(x) \prod_{j=1}^n x_j^{2\lambda_j} dx \\ &= \int_{(0, \infty)^n} \sum_{j=1}^n y_j^2 \int_0^\infty e^{-t \sum_{j=1}^n y_j^2} \phi(t) dt h_{\lambda_1, \dots, \lambda_n}(f)(y) h_{\lambda_1, \dots, \lambda_n}(g)(y) \prod_{j=1}^n y_j^{2\lambda_j} dy \\ &= \int_0^\infty \phi(t) \int_{(0, \infty)^n} \sum_{j=1}^n y_j^2 e^{-t \sum_{j=1}^n y_j^2} h_{\lambda_1, \dots, \lambda_n}(f)(y) h_{\lambda_1, \dots, \lambda_n}(g)(y) \prod_{j=1}^n y_j^{2\lambda_j} dy dt. \end{aligned}$$

The interchange of the order of integration is justify because we can write

$$\begin{aligned} & \int_{(0, \infty)^n} \sum_{j=1}^n y_j^2 \int_0^\infty |\phi(t)| e^{-t \sum_{j=1}^n y_j^2} dt |h_{\lambda_1, \dots, \lambda_n}(f)(y)| |h_{\lambda_1, \dots, \lambda_n}(g)(y)| \prod_{j=1}^n y_j^{2\lambda_j} dy \\ & \leq C \int_{(0, \infty)^n} |h_{\lambda_1, \dots, \lambda_n}(f)(y)| |h_{\lambda_1, \dots, \lambda_n}(g)(y)| \prod_{j=1}^n y_j^{2\lambda_j} dy \\ & \leq C \|h_{\lambda_1, \dots, \lambda_n}(f)\|_{L^2((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)} \|h_{\lambda_1, \dots, \lambda_n}(g)\|_{L^2((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)} \\ & = C \|f\|_{L^2((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)} \|g\|_{L^2((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)} < \infty, \end{aligned}$$

where we have used that $h_{\lambda_1, \dots, \lambda_n}$ is an isometry in $L^2((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$.

Then, by using again (12), we obtain

$$\begin{aligned} & \int_{(0,\infty)^n} T_{\lambda_1,\dots,\lambda_n}^m(f)(x)g(x) \prod_{j=1}^n x_j^{2\lambda_j} dx \\ &= \int_0^\infty \phi(t) \int_{(0,\infty)^n} h_{\lambda_1,\dots,\lambda_n} \left(\sum_{j=1}^n y_j^2 e^{-t \sum_{j=1}^n y_j^2} h_{\lambda_1,\dots,\lambda_n}(f)(y) \right) (x) g(x) \prod_{j=1}^n x_j^{2\lambda_j} dx dt. \end{aligned}$$

According to (7) and (8), for each $0 \leq r \leq n$,

$$\begin{aligned} & \int_0^{1/x_1} \cdots \int_0^{1/x_r} \int_{1/x_{r+1}}^\infty \cdots \int_{1/x_n}^\infty \left| \prod_{j=1}^n (x_j y_j)^{-\lambda_j+1/2} J_{\lambda_j-1/2}(x_j y_j) \right| \\ & \times \sum_{j=1}^n y_j^2 e^{-t \sum_{j=1}^n y_j^2} |h_{\lambda_1,\dots,\lambda_n}(f)(y)| \prod_{j=1}^n y_j^{2\lambda_j} dy \\ & \leq C \prod_{j=r+1}^n x_j^{-\lambda_j} \int_{(0,\infty)^n} \sum_{j=1}^n y_j^2 e^{-t \sum_{j=1}^n y_j^2} |h_{\lambda_1,\dots,\lambda_n}(f)(y)| \prod_{j=1}^r y_j^{2\lambda_j} \prod_{j=r+1}^n y_j^{\lambda_j} dy, \quad x \in (0,\infty)^n. \end{aligned}$$

Then, since $|y|^l h_{\lambda_1,\dots,\lambda_n}(f)$ is bounded on $(0,\infty)^n$, for every $l \in \mathbb{N}$, and $g \in C_c^\infty((0,\infty)^n)$, we get, for every $0 \leq r \leq n$,

$$\begin{aligned} & \int_0^\infty |\phi(t)| \int_{(0,\infty)^n} \int_0^{1/x_1} \cdots \int_0^{1/x_r} \int_{1/x_{r+1}}^\infty \cdots \int_{1/x_n}^\infty \left| \prod_{j=1}^n (x_j y_j)^{-\lambda_j+1/2} J_{\lambda_j-1/2}(x_j y_j) \right| \\ & \times \sum_{j=1}^n y_j^2 e^{-t \sum_{j=1}^n y_j^2} |h_{\lambda_1,\dots,\lambda_n}(f)(y)| \prod_{j=1}^n y_j^{2\lambda_j} dy |g(x)| \prod_{j=1}^n x_j^{2\lambda_j} dx dt \\ & \leq C \int_0^\infty \int_{(0,\infty)^n} \int_{(0,\infty)^n} \sum_{j=1}^n y_j^2 e^{-t \sum_{j=1}^n y_j^2} |h_{\lambda_1,\dots,\lambda_n}(f)(y)| |g(x)| \\ & \times \prod_{j=1}^r y_j^{2\lambda_j} \prod_{j=r+1}^n y_j^{\lambda_j} \prod_{j=1}^r x_j^{2\lambda_j} \prod_{j=r+1}^n x_j^{\lambda_j} dy dx dt \\ & \leq C \int_{(0,\infty)^n} |g(x)| \prod_{j=1}^r x_j^{2\lambda_j} \prod_{j=r+1}^n x_j^{\lambda_j} dx \\ & \times \int_{(0,\infty)^n} \left(\int_0^\infty \sum_{j=1}^n y_j^2 e^{-t \sum_{j=1}^n y_j^2} dt \right) |h_{\lambda_1,\dots,\lambda_n}(f)(y)| \prod_{j=1}^r y_j^{2\lambda_j} \prod_{j=r+1}^n y_j^{\lambda_j} dy \\ & \leq C \left(\int_{(0,\infty)^n} |g(x)| \prod_{j=1}^r x_j^{2\lambda_j} \prod_{j=r+1}^n x_j^{\lambda_j} dx \right) \left(\int_{(0,\infty)^n} |h_{\lambda_1,\dots,\lambda_n}(f)(y)| \prod_{j=1}^r y_j^{2\lambda_j} \prod_{j=r+1}^n y_j^{\lambda_j} dy \right) < \infty. \end{aligned}$$

Note that these estimates also show that the function

$$G(x) = \int_0^\infty \phi(t) h_{\lambda_1,\dots,\lambda_n} \left(\sum_{j=1}^n y_j^2 e^{-t \sum_{j=1}^n y_j^2} h_{\lambda_1,\dots,\lambda_n}(f)(y) \right) (x) dt, \quad x \in (0,\infty)^n,$$

is in $L_{loc}^1((0,\infty)^n)$.

We conclude that

$$\begin{aligned} & \int_{(0,\infty)^n} T_{\lambda_1,\dots,\lambda_n}^m(f)(x)g(x) \prod_{j=1}^n x_j^{2\lambda_j} dx \\ &= \int_{(0,\infty)^n} \left\{ \int_0^\infty \phi(t) h_{\lambda_1,\dots,\lambda_n} \left(\sum_{j=1}^n y_j^2 e^{-t \sum_{j=1}^n y_j^2} h_{\lambda_1,\dots,\lambda_n}(f)(y) \right) (x) dt \right\} g(x) \prod_{j=1}^n x_j^{2\lambda_j} dx. \end{aligned}$$

Thus, the proof of this Lemma finishes because $T_{\lambda_1, \dots, \lambda_n}^m(f) \in L^2((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx) \subset L_{loc}^1((0, \infty)^n)$. $\square$

We now establish a crucial result in order to prove Theorem 1.1.

Proposition 2.1. *Let $\lambda_j > -1/2$, $j = 1, \dots, n$. Assume that $f \in C_c^\infty((0, \infty)^n)$. Then*

$$T_{\lambda_1, \dots, \lambda_n}^m(f)(x) = - \lim_{\varepsilon \rightarrow 0^+} \left(\alpha(\varepsilon) f(x) + \int_{(0, \infty)^n, |y-x| > \varepsilon} f(y) K_{\lambda_1, \dots, \lambda_n}^\phi(x, y) \prod_{j=1}^n y_j^{2\lambda_j} dy \right), \text{ a.e. } x \in (0, \infty)^n,$$

where

$$K_{\lambda_1, \dots, \lambda_n}^\phi(x, y) = \int_0^\infty \phi(t) \frac{\partial}{\partial t} \prod_{j=1}^n W_t^{\lambda_j}(x_j, y_j) dt, \quad x, y \in (0, \infty)^n, \quad x \neq y,$$

and α is a bounded function on $(0, \infty)$. Moreover, if there exists the limit $\phi(0^+) = \lim_{t \rightarrow 0^+} \phi(t)$, then

$$T_{\lambda_1, \dots, \lambda_n}^m(f)(x) = C\phi(0^+)f(x) - \lim_{\varepsilon \rightarrow 0^+} \left(\int_{(0, \infty)^n, |y-x| > \varepsilon} f(y) K_{\lambda_1, \dots, \lambda_n}^\phi(x, y) \prod_{j=1}^n y_j^{2\lambda_j} dy \right), \text{ a.e. } x \in (0, \infty)^n,$$

being C a positive constant.

Proof. Assume that $n \geq 2$. When $n = 1$ we can proceed in a similar way. By Lemma 2.1 and (11) we can write, for a.e. $x \in (0, \infty)^n$,

$$(13) \quad T_{\lambda_1, \dots, \lambda_n}^m(f)(x) = \int_0^\infty \phi(t) h_{\lambda_1, \dots, \lambda_n} \left(e^{-t \sum_{j=1}^n y_j^2} h_{\lambda_1, \dots, \lambda_n}(\Delta_{\lambda_1, \dots, \lambda_n} f)(y) \right) (x) dt.$$

Fix $x \in (0, \infty)^n$ such that (13) holds. According to (7) and (8), for each $0 \leq r, s \leq n$, it follows that,

$$\begin{aligned} & \int_0^{1/x_1} \cdots \int_0^{1/x_r} \int_{1/x_{r+1}}^\infty \cdots \int_{1/x_n}^\infty \prod_{j=1}^n \left| (x_j y_j)^{-\lambda_j+1/2} J_{\lambda_j-1/2}(x_j y_j) \right| e^{-t \sum_{j=1}^n y_j^2} \\ & \quad \times \int_0^{1/y_1} \cdots \int_0^{1/y_s} \int_{1/y_{s+1}}^\infty \cdots \int_{1/y_n}^\infty \prod_{j=1}^n \left| (y_j z_j)^{-\lambda_j+1/2} J_{\lambda_j-1/2}(y_j z_j) \right| \\ & \quad \times |\Delta_{\lambda_1, \dots, \lambda_n} f(z)| \prod_{j=1}^n z_j^{2\lambda_j} dz \prod_{j=1}^n y_j^{2\lambda_j} dy \\ & \leq C \prod_{j=r+1}^n x_j^{-\lambda_j} \int_{(0, \infty)^n} e^{-t \sum_{j=1}^n y_j^2} \prod_{j=r+1}^n y_j^{-\lambda_j} \prod_{j=1}^s y_j^{2\lambda_j} \prod_{j=s+1}^n y_j^{\lambda_j} dy \\ & \quad \times \int_{(0, \infty)^n} |\Delta_{\lambda_1, \dots, \lambda_n} f(z)| \prod_{j=1}^s z_j^{2\lambda_j} \prod_{j=s+1}^n z_j^{\lambda_j} dz < \infty, \quad t > 0. \end{aligned}$$

Then, by interchanging the order of integration and by using (10), we get

$$\begin{aligned} & h_{\lambda_1, \dots, \lambda_n} \left(e^{-t \sum_{j=1}^n y_j^2} h_{\lambda_1, \dots, \lambda_n}(\Delta_{\lambda_1, \dots, \lambda_n} f)(y) \right) (x) \\ & = \int_{(0, \infty)^n} \Delta_{\lambda_1, \dots, \lambda_n} f(z) \prod_{j=1}^n \int_0^\infty e^{-t y_j^2} (x_j y_j)^{-\lambda_j+1/2} J_{\lambda_j-1/2}(x_j y_j) \\ & \quad \times (y_j z_j)^{-\lambda_j+1/2} J_{\lambda_j-1/2}(y_j z_j) y_j^{2\lambda_j} dy_j \prod_{j=1}^n z_j^{2\lambda_j} dz \\ (14) \quad & = \int_{(0, \infty)^n} \Delta_{\lambda_1, \dots, \lambda_n} f(z) \prod_{j=1}^n W_t^{\lambda_j}(x_j, z_j) \prod_{j=1}^n z_j^{2\lambda_j} dz, \quad t > 0, \end{aligned}$$

where, for every $\alpha > -1/2$, $W_t^\alpha(u, v)$, $t, u, v \in (0, \infty)$, is defined in (3).

We choose $a > 1$ such that $\text{supp } f \subset K^n$, where $K = [1/a, a]$. If $\alpha > -1/2$, by (4), (5) and (6), we obtain, for every $b > 0$,

$$\begin{aligned}
\left| W_t^\alpha(u, v) - \frac{t^{-\alpha-1/2}}{2^{2\alpha}\Gamma(\alpha+1/2)} \right| &= \left| \frac{1}{(2t)^{\alpha+1/2}} \left(\frac{uv}{2t} \right)^{-\alpha+1/2} I_{\alpha-1/2} \left(\frac{uv}{2t} \right) e^{-\frac{u^2+v^2}{4t}} - \frac{t^{-\alpha-1/2}}{2^{2\alpha}\Gamma(\alpha+1/2)} \right| \\
&\leq \frac{1}{(2t)^{\alpha+1/2}} e^{-\frac{u^2+v^2}{4t}} \left| \left(\frac{uv}{2t} \right)^{-\alpha+1/2} I_{\alpha-1/2} \left(\frac{uv}{2t} \right) - \frac{1}{2^{\alpha-1/2}\Gamma(\alpha+1/2)} \right| \\
&\quad + \frac{t^{-\alpha-1/2}}{2^{2\alpha}\Gamma(\alpha+1/2)} \left| e^{-\frac{u^2+v^2}{4t}} - 1 \right| \\
&\leq \frac{uv}{(2t)^{\alpha+3/2}} e^{-\frac{u^2+v^2}{4t}} \sup_{z \in (0, \frac{uv}{2t})} \left| \frac{d}{dz} \left(z^{-\alpha+1/2} I_{\alpha-1/2}(z) \right) \right| + C \frac{u^2+v^2}{t^{\alpha+3/2}} \\
&\leq C \left(\frac{(uv)^2}{t^{\alpha+5/2}} + \frac{u^2+v^2}{t^{\alpha+3/2}} \right) \\
&\leq C \frac{1}{t^{\alpha+3/2}}, \quad t > 1, \quad 0 < u \leq b, \text{ and } v \in K.
\end{aligned}$$

Then, by using again (4) and (5), it follows that

$$\begin{aligned}
&\left| \prod_{j=1}^n W_t^{\lambda_j}(x_j, y_j) - \frac{t^{-\sum_{j=1}^n (\lambda_j+1/2)}}{\prod_{j=1}^n 2^{2\lambda_j}\Gamma(\lambda_j+1/2)} \right| \\
&\leq \sum_{i=1}^n \prod_{j=1}^{i-1} \frac{t^{-(\lambda_j+1/2)}}{2^{2\lambda_j}\Gamma(\lambda_j+1/2)} \left| W_t^{\lambda_i}(x_i, y_i) - \frac{t^{-(\lambda_i+1/2)}}{2^{2\lambda_i}\Gamma(\lambda_i+1/2)} \right| \prod_{j=i+1}^n W_t^{\lambda_j}(x_j, y_j) \\
(15) \quad &\leq C \frac{1}{t^{\sum_{j=1}^n (\lambda_j+1/2)+1}}, \quad t > 1 \text{ and } y \in K^n.
\end{aligned}$$

On the other hand, by (7) and (11) we have

$$\begin{aligned}
&\int_{(0, \infty)^n} \Delta_{\lambda_1, \dots, \lambda_n} f(z) \prod_{j=1}^n z_j^{2\lambda_j} dz = \prod_{j=1}^n 2^{\lambda_j-1/2} \Gamma(\lambda_j+1/2) \\
&\quad \times \lim_{y \rightarrow 0} \int_{(0, \infty)^n} \Delta_{\lambda_1, \dots, \lambda_n} f(z) \prod_{j=1}^n (z_j y_j)^{-\lambda_j+1/2} J_{\lambda_j-1/2}(z_j y_j) \prod_{j=1}^n z_j^{2\lambda_j} dz \\
(16) \quad &= \prod_{j=1}^n 2^{\lambda_j-1/2} \Gamma(\lambda_j+1/2) \lim_{y \rightarrow 0} \sum_{j=1}^n y_j^2 h_{\lambda_1, \dots, \lambda_n}(f)(y) = 0.
\end{aligned}$$

According to (13), (14), and (16) (suggested by (15)), we can write

$$\begin{aligned}
T_{\lambda_1, \dots, \lambda_n}^m(f)(x) &= \int_0^\infty \phi(t) \int_{(0, \infty)^n} \Delta_{\lambda_1, \dots, \lambda_n} f(z) \left(\prod_{j=1}^n W_t^{\lambda_j}(x_j, z_j) - \frac{\chi_{(1, \infty)}(t) t^{-\sum_{j=1}^n (\lambda_j+1/2)}}{\prod_{j=1}^n 2^{2\lambda_j}\Gamma(\lambda_j+1/2)} \right) \\
(17) \quad &\quad \times \prod_{j=1}^n z_j^{2\lambda_j} dz dt.
\end{aligned}$$

The last integral is absolutely convergent. Indeed, in order to do this we split the integral in the following way

$$\begin{aligned}
& \int_0^\infty |\phi(t)| \int_{(0,\infty)^n} |\Delta_{\lambda_1, \dots, \lambda_n} f(z)| \left| \prod_{j=1}^n W_t^{\lambda_j}(x_j, z_j) - \frac{\chi_{(1,\infty)}(t)t^{-\sum_{j=1}^n (\lambda_j+1/2)}}{\prod_{j=1}^n 2^{2\lambda_j} \Gamma(\lambda_j + 1/2)} \right| \prod_{j=1}^n z_j^{2\lambda_j} dz dt \\
& \leq C \left(\int_0^1 + \int_1^\infty \right) \int_{(0,\infty)^n} |\Delta_{\lambda_1, \dots, \lambda_n} f(z)| \left| \prod_{j=1}^n W_t^{\lambda_j}(x_j, z_j) - \frac{\chi_{(1,\infty)}(t)t^{-\sum_{j=1}^n (\lambda_j+1/2)}}{\prod_{j=1}^n 2^{2\lambda_j} \Gamma(\lambda_j + 1/2)} \right| \prod_{j=1}^n z_j^{2\lambda_j} dz dt \\
& = I_1(x) + I_2(x).
\end{aligned}$$

Since by using the inversion formula for Hankel transform and (10) we get

$$\int_0^\infty W_t^\alpha(u, z) z^{2\alpha} dz = 1, \quad u \in (0, \infty),$$

when $\alpha > -1/2$, it follows that

$$I_1(x) \leq C \int_0^1 \int_{(0,\infty)^n} |\Delta_{\lambda_1, \dots, \lambda_n} f(z)| \prod_{j=1}^n W_t^{\lambda_j}(x_j, z_j) \prod_{j=1}^n z_j^{2\lambda_j} dz dt \leq C.$$

Also, by (15) we obtain

$$I_2(x) \leq C \int_1^\infty \frac{1}{\sum_{j=1}^n (\lambda_j + 1/2) + 1} \int_{K^n} |\Delta_{\lambda_1, \dots, \lambda_n} f(z)| \prod_{j=1}^n z_j^{2\lambda_j} dz dt \leq C.$$

By (17) we have

$$\begin{aligned}
(18) \quad T_{\lambda_1, \dots, \lambda_n}^m(f)(x) &= \lim_{\varepsilon \rightarrow 0^+} \int_0^\infty \phi(t) \int_{\substack{(0, \infty)^n \\ |z-x| > \varepsilon}} \Delta_{\lambda_1, \dots, \lambda_n} f(z) \\
&\quad \times \left(\prod_{j=1}^n W_t^{\lambda_j}(x_j, z_j) - \frac{\chi_{(1,\infty)}(t)t^{-\sum_{j=1}^n (\lambda_j+1/2)}}{\prod_{j=1}^n 2^{2\lambda_j} \Gamma(\lambda_j + 1/2)} \right) \prod_{j=1}^n z_j^{2\lambda_j} dz dt.
\end{aligned}$$

Assume that ε is small enough, for instance, $0 < \varepsilon < x_i/2$, $i = 1, \dots, n$. We now analyze the integral

$$I^\varepsilon(x, t) = \int_{\substack{K^n \\ |z-x| > \varepsilon}} \Delta_{\lambda_1, z_1} f(z) \left(\prod_{j=1}^n W_t^{\lambda_j}(x_j, z_j) - \frac{\chi_{(1,\infty)}(t)t^{-\sum_{j=1}^n (\lambda_j+1/2)}}{\prod_{j=1}^n 2^{2\lambda_j} \Gamma(\lambda_j + 1/2)} \right) \prod_{j=1}^n z_j^{2\lambda_j} dz, \quad t > 0.$$

The study of the integral involving Δ_{λ_j, z_j} , $j = 2, \dots, n$, can be made in a similar way. We write $\bar{y} = (y_2, \dots, y_n)$ when $y = (y_1, y_2, \dots, y_n) \in (0, \infty)^n$. We can write

$$\begin{aligned}
(19) \quad I^\varepsilon(x, t) &= \int_{|\bar{z}-\bar{x}| > \varepsilon} \int_0^\infty \Delta_{\lambda_1, z_1} f(z) \left(\prod_{j=1}^n W_t^{\lambda_j}(x_j, z_j) - \frac{\chi_{(1,\infty)}(t)t^{-\sum_{j=1}^n (\lambda_j+1/2)}}{\prod_{j=1}^n 2^{2\lambda_j} \Gamma(\lambda_j + 1/2)} \right) \prod_{j=1}^n z_j^{2\lambda_j} dz \\
&\quad + \int_{|\bar{z}-\bar{x}| < \varepsilon} \left(\int_0^{x_1 - \sqrt{\varepsilon^2 - |\bar{z}-\bar{x}|^2}} + \int_{x_1 + \sqrt{\varepsilon^2 - |\bar{z}-\bar{x}|^2}}^\infty \right) \Delta_{\lambda_1, z_1} f(z) \\
&\quad \times \left(\prod_{j=1}^n W_t^{\lambda_j}(x_j, z_j) - \frac{\chi_{(1,\infty)}(t)t^{-\sum_{j=1}^n (\lambda_j+1/2)}}{\prod_{j=1}^n 2^{2\lambda_j} \Gamma(\lambda_j + 1/2)} \right) \prod_{j=1}^n z_j^{2\lambda_j} dz \\
&= I_1^\varepsilon(x, t) + I_2^\varepsilon(x, t), \quad t > 0.
\end{aligned}$$

By partial integration we obtain

$$\begin{aligned}
& \int_0^\infty \Delta_{\lambda_1, z_1} f(z) \left(\prod_{j=1}^n W_t^{\lambda_j}(x_j, z_j) - \frac{\chi_{(1, \infty)}(t) t^{-\sum_{j=1}^n (\lambda_j + 1/2)}}{\prod_{j=1}^n 2^{2\lambda_j} \Gamma(\lambda_j + 1/2)} \right) z_1^{2\lambda_1} dz_1 \\
&= -z_1^{2\lambda_1} \frac{\partial}{\partial z_1} f(z) \left(\prod_{j=1}^n W_t^{\lambda_j}(x_j, z_j) - \frac{\chi_{(1, \infty)}(t) t^{-\sum_{j=1}^n (\lambda_j + 1/2)}}{\prod_{j=1}^n 2^{2\lambda_j} \Gamma(\lambda_j + 1/2)} \right) \Big|_0^\infty \\
&\quad + z_1^{2\lambda_1} \frac{\partial}{\partial z_1} \left(\prod_{j=1}^n W_t^{\lambda_j}(x_j, z_j) \right) f(z) \Big|_0^\infty + \int_0^\infty f(z) \Delta_{\lambda_1, z_1} W_t^{\lambda_1}(x_1, z_1) \prod_{j=2}^n W_t^{\lambda_j}(x_j, z_j) z_1^{2\lambda_1} dz_1 \\
&= \int_{1/a}^a f(z) \Delta_{\lambda_1, z_1} W_t^{\lambda_1}(x_1, z_1) \prod_{j=2}^n W_t^{\lambda_j}(x_j, z_j) z_1^{2\lambda_1} dz_1, \quad t > 0 \text{ and } \bar{z} \in K^{n-1}.
\end{aligned}$$

Differentiating in (10), by using (9) and [11, (5.3.6), p. 103], we get $-\Delta_{\alpha, u} W_t^\alpha(u, v) = \frac{\partial}{\partial t} W_t^\alpha(u, v)$, $u, v, t > 0$, $\alpha > -1/2$, and it follows that

$$\begin{aligned}
I_1^\varepsilon(x, t) &= \int_{|\bar{z}-\bar{x}|>\varepsilon} \int_0^\infty \Delta_{\lambda_1, z_1} f(z) \left(\prod_{j=1}^n W_t^{\lambda_j}(x_j, z_j) - \frac{\chi_{(1, \infty)}(t) t^{-\sum_{j=1}^n (\lambda_j + 1/2)}}{\prod_{j=1}^n 2^{2\lambda_j} \Gamma(\lambda_j + 1/2)} \right) \prod_{j=1}^n z_j^{2\lambda_j} dz \\
(20) \quad &= - \int_{|\bar{z}-\bar{x}|>\varepsilon} \int_0^\infty f(z) \frac{\partial}{\partial t} W_t^{\lambda_1}(x_1, z_1) \prod_{j=2}^n W_t^{\lambda_j}(x_j, z_j) \prod_{j=1}^n z_j^{2\lambda_j} dz, \quad t > 0.
\end{aligned}$$

In a similar way we obtain

$$\begin{aligned}
& \left(\int_0^{x_1 - \sqrt{\varepsilon^2 - |\bar{z} - \bar{x}|^2}} + \int_{x_1 + \sqrt{\varepsilon^2 - |\bar{z} - \bar{x}|^2}}^\infty \right) \Delta_{\lambda_1, z_1} f(z) \left(\prod_{j=1}^n W_t^{\lambda_j}(x_j, z_j) - \frac{\chi_{(1, \infty)}(t) t^{-\sum_{j=1}^n (\lambda_j + 1/2)}}{\prod_{j=1}^n 2^{2\lambda_j} \Gamma(\lambda_j + 1/2)} \right) z_1^{2\lambda_1} dz_1 \\
&= \left(\int_0^{x_1 - \sqrt{\varepsilon^2 - |\bar{z} - \bar{x}|^2}} + \int_{x_1 + \sqrt{\varepsilon^2 - |\bar{z} - \bar{x}|^2}}^\infty \right) f(z) \Delta_{\lambda_1, z_1} W_t^{\lambda_1}(x_1, z_1) \prod_{j=2}^n W_t^{\lambda_j}(x_j, z_j) z_1^{2\lambda_1} dz_1 \\
&\quad - \left(H_1(x, x_1 - \sqrt{\varepsilon^2 - |\bar{z} - \bar{x}|^2}, \bar{z}, t) - H_1(x, x_1 + \sqrt{\varepsilon^2 - |\bar{z} - \bar{x}|^2}, \bar{z}, t) \right) \\
&\quad + H_2(x, x_1 - \sqrt{\varepsilon^2 - |\bar{z} - \bar{x}|^2}, \bar{z}, t) - H_2(x, x_1 + \sqrt{\varepsilon^2 - |\bar{z} - \bar{x}|^2}, \bar{z}, t), \quad |\bar{x} - \bar{z}| < \varepsilon \text{ and } t > 0.
\end{aligned}$$

where

$$H_1(x, z, t) = z_1^{2\lambda_1} \frac{\partial}{\partial z_1} f(z) \left(\prod_{j=1}^n W_t^{\lambda_j}(x_j, z_j) - \frac{\chi_{(1, \infty)}(t) t^{-\sum_{j=1}^n (\lambda_j + 1/2)}}{\prod_{j=1}^n 2^{2\lambda_j} \Gamma(\lambda_j + 1/2)} \right), \quad t > 0 \text{ and } z \in (0, \infty)^n,$$

and

$$H_2(x, z, t) = z_1^{2\lambda_1} \frac{\partial}{\partial z_1} \left(\prod_{j=1}^n W_t^{\lambda_j}(x_j, z_j) \right) f(z), \quad t > 0 \text{ and } z \in (0, \infty)^n.$$

We have, by (20), that

$$\begin{aligned}
I_1^\varepsilon(x, t) &+ \int_{|\bar{z}-\bar{x}|<\varepsilon} \left(\int_0^{x_1 - \sqrt{\varepsilon^2 - |\bar{z} - \bar{x}|^2}} + \int_{x_1 + \sqrt{\varepsilon^2 - |\bar{z} - \bar{x}|^2}}^\infty \right) f(z) \Delta_{\lambda_1, z_1} W_t^{\lambda_1}(x_1, z_1) \prod_{j=2}^n W_t^{\lambda_j}(x_j, z_j) \prod_{j=1}^n z_j^{2\lambda_j} dz \\
(21) \quad &= - \int_{|z-x|>\varepsilon} f(z) \frac{\partial}{\partial t} W_t^{\lambda_1}(x_1, z_1) \prod_{j=2}^n W_t^{\lambda_j}(x_j, z_j) \prod_{j=1}^n z_j^{2\lambda_j} dz, \quad t > 0.
\end{aligned}$$

Moreover, according to (4), (5), and (6) it follows that, if $\mathbb{K}$ is a compact subset of $(0, \infty)$, for every $z_1 \in \mathbb{K}$,

$$\begin{aligned}
 \left| \frac{\partial}{\partial z_1} W_t^{\lambda_1}(x_1, z_1) \right| &= \left| \frac{\partial}{\partial z_1} \left(\frac{1}{(2t)^{\lambda_1+1/2}} \left(\frac{x_1 z_1}{2t} \right)^{-\lambda_1+1/2} I_{\lambda_1-1/2} \left(\frac{x_1 z_1}{2t} \right) e^{-(x_1^2+z_1^2)/4t} \right) \right| \\
 &= \frac{1}{(2t)^{\lambda_1+1/2}} \left| \frac{x_1}{2t} \left(\frac{x_1 z_1}{2t} \right)^{-\lambda_1+1/2} I_{\lambda_1+1/2} \left(\frac{x_1 z_1}{2t} \right) \right. \\
 &\quad \left. - \frac{z_1}{2t} \left(\frac{x_1 z_1}{2t} \right)^{-\lambda_1+1/2} I_{\lambda_1-1/2} \left(\frac{x_1 z_1}{2t} \right) \right| e^{-(x_1^2+z_1^2)/4t} \\
 (22) \quad &\leq C \begin{cases} \frac{1}{t^{\lambda_1+3/2}} & , \quad t \geq 1 \\ \frac{1}{t} e^{-(x_1-z_1)^2/8t} & , \quad 0 < t < 1. \end{cases}
 \end{aligned}$$

Then, since $f \in C_c^\infty((0, \infty)^n)$, by using main value theorem, (4), (15) and (22) imply that

$$\begin{aligned}
 &|H_1(x, x_1 - \sqrt{\varepsilon^2 - |\bar{z} - \bar{x}|^2}, \bar{z}, t) - H_1(x, x_1 + \sqrt{\varepsilon^2 - |\bar{z} - \bar{x}|^2}, \bar{z}, t)| \\
 &\leq |(x_1 - \sqrt{\varepsilon^2 - |\bar{z} - \bar{x}|^2})^{2\lambda_1} \left(\frac{\partial}{\partial z_1} f \right)(x_1 - \sqrt{\varepsilon^2 - |\bar{z} - \bar{x}|^2}, \bar{z}) \\
 &\quad - (x_1 + \sqrt{\varepsilon^2 - |\bar{z} - \bar{x}|^2})^{2\lambda_1} \left(\frac{\partial}{\partial z_1} f \right)(x_1 + \sqrt{\varepsilon^2 - |\bar{z} - \bar{x}|^2}, \bar{z})| \\
 &\quad \times \left| W_t^{\lambda_1}(x_1, x_1 + \sqrt{\varepsilon^2 - |\bar{z} - \bar{x}|^2}) \prod_{j=2}^n W_t^{\lambda_j}(x_j, z_j) - \frac{\chi_{(1,\infty)}(t) t^{-\sum_{j=1}^n (\lambda_j+1/2)}}{\prod_{j=1}^n 2^{2\lambda_j} \Gamma(\lambda_j + 1/2)} \right| \\
 &\quad + \left| (x_1 - \sqrt{\varepsilon^2 - |\bar{z} - \bar{x}|^2})^{2\lambda_1} \left(\frac{\partial}{\partial z_1} f \right)(x_1 - \sqrt{\varepsilon^2 - |\bar{z} - \bar{x}|^2}, \bar{z}) \right| \\
 &\quad \times \left| W_t^{\lambda_1}(x_1, x_1 - \sqrt{\varepsilon^2 - |\bar{z} - \bar{x}|^2}) - W_t^{\lambda_1}(x_1, x_1 + \sqrt{\varepsilon^2 - |\bar{z} - \bar{x}|^2}) \right| \prod_{j=2}^n W_t^{\lambda_j}(x_j, z_j) \\
 (23) \quad &\leq C \frac{\varepsilon}{\sum_{j=1}^n (\lambda_j+1/2)+1}, \quad t \geq 1, \quad \bar{z} \in K^{n-1} \quad \text{and} \quad |\bar{z} - \bar{x}| < \varepsilon.
 \end{aligned}$$

Also, from (5) and (22) we deduce

$$\begin{aligned}
 &|H_1(x, x_1 - \sqrt{\varepsilon^2 - |\bar{z} - \bar{x}|^2}, \bar{z}, t) - H_1(x, x_1 + \sqrt{\varepsilon^2 - |\bar{z} - \bar{x}|^2}, \bar{z}, t)| \\
 &\leq C \left\{ \sqrt{\varepsilon^2 - |\bar{z} - \bar{x}|^2} \frac{(x_1(x_1 + \sqrt{\varepsilon^2 - |\bar{z} - \bar{x}|^2}))^{-\lambda_1}}{\sqrt{t}} \right. \\
 &\quad \left. + \left| (x_1 - \sqrt{\varepsilon^2 - |\bar{z} - \bar{x}|^2})^{2\lambda_1} \left(\frac{\partial}{\partial z_1} f \right)(x_1 - \sqrt{\varepsilon^2 - |\bar{z} - \bar{x}|^2}, \bar{z}) \right| \right. \\
 &\quad \left. \times \int_{x_1 - \sqrt{\varepsilon^2 - |\bar{z} - \bar{x}|^2}}^{x_1 + \sqrt{\varepsilon^2 - |\bar{z} - \bar{x}|^2}} \left| \frac{\partial}{\partial z_1} W_t^{\lambda_1}(x_1, z_1) \right| dz_1 \right\} \prod_{j=2}^n W_t^{\lambda_j}(x_j, z_j) \\
 &\leq C \left\{ \frac{\varepsilon}{\sqrt{t}} + \int_{x_1 - \sqrt{\varepsilon^2 - |\bar{z} - \bar{x}|^2}}^{x_1 + \sqrt{\varepsilon^2 - |\bar{z} - \bar{x}|^2}} \frac{e^{-|x_1 - z_1|^2/8t}}{t} dz_1 \right\} \\
 (24) \quad &\times \prod_{j=2}^n \frac{1}{\sqrt{t}} e^{-(x_j - z_j)^2/4t}, \quad 0 < t < 1, \quad \bar{z} \in K^{n-1} \quad \text{and} \quad |\bar{z} - \bar{x}| < \varepsilon.
 \end{aligned}$$

By (23), we can write

$$\begin{aligned}
 & \int_1^\infty |\phi(t)| \int_{\bar{z} \in K^{n-1}, |\bar{z}-\bar{x}| < \varepsilon} |H_1(x, x_1 - \sqrt{\varepsilon^2 - |\bar{z}-\bar{x}|^2}, \bar{z}, t) - H_1(x, x_1 + \sqrt{\varepsilon^2 - |\bar{z}-\bar{x}|^2}, \bar{z}, t)| \prod_{j=2}^n z_j^{2\lambda_j} d\bar{z} dt \\
 (25) \quad & \leq C\varepsilon \int_1^\infty \int_{|\bar{z}-\bar{x}| < \varepsilon} \frac{1}{\sum_{j=2}^n (\lambda_j + 1/2) + 1} \prod_{j=2}^n z_j^{2\lambda_j} d\bar{z} dt \longrightarrow 0, \quad \varepsilon \rightarrow 0^+.
 \end{aligned}$$

By (24) and by using [20, Lemma 1.1], when $n > 2$ it follows

$$\begin{aligned}
 & \int_0^1 |\phi(t)| \int_{\bar{z} \in K^{n-1}, |\bar{z}-\bar{x}| < \varepsilon} |H_1(x, x_1 - \sqrt{\varepsilon^2 - |\bar{z}-\bar{x}|^2}, \bar{z}, t) - H_1(x, x_1 + \sqrt{\varepsilon^2 - |\bar{z}-\bar{x}|^2}, \bar{z}, t)| \prod_{j=2}^n z_j^{2\lambda_j} d\bar{z} dt \\
 (26) \quad & \leq C \left(\varepsilon \int_0^1 \int_{\bar{z} \in K^{n-1}, |\bar{z}-\bar{x}| < \varepsilon} \frac{e^{-|\bar{z}-\bar{x}|^2/4t}}{t^{n/2}} d\bar{z} dt + \int_0^1 \int_{\bar{z} \in K^{n-1}, |\bar{z}-\bar{x}| < \varepsilon} \int_{x_1 - \sqrt{\varepsilon^2 - |\bar{z}-\bar{x}|^2}}^{x_1 + \sqrt{\varepsilon^2 - |\bar{z}-\bar{x}|^2}} \frac{e^{-|z-x|^2/8t}}{t^{(n+1)/2}} dz dt \right) \\
 & \leq C \left(\varepsilon \int_{|\bar{z}-\bar{x}| < \varepsilon} \frac{d\bar{z}}{|\bar{z}-\bar{x}|^{n-2}} + \int_{|z-x| < \varepsilon} \frac{dz}{|z-x|^{n-1}} \right) \rightarrow 0, \quad \text{as } \varepsilon \rightarrow 0^+.
 \end{aligned}$$

For $n = 2$ we can proceed analogously.

We now write, for each $t > 0$, $\bar{z} \in K^{n-1}$ and $|\bar{z}-\bar{x}| < \varepsilon$,

$$\begin{aligned}
 & |H_2(x, x_1 - \sqrt{\varepsilon^2 - |\bar{z}-\bar{x}|^2}, \bar{z}, t) - H_2(x, x_1 + \sqrt{\varepsilon^2 - |\bar{z}-\bar{x}|^2}, \bar{z}, t)| \\
 & \leq |(x_1 - \sqrt{\varepsilon^2 - |\bar{z}-\bar{x}|^2})^{2\lambda_1} f(x_1 - \sqrt{\varepsilon^2 - |\bar{z}-\bar{x}|^2}, \bar{z}) - (x_1 + \sqrt{\varepsilon^2 - |\bar{z}-\bar{x}|^2})^{2\lambda_1} f(x_1 + \sqrt{\varepsilon^2 - |\bar{z}-\bar{x}|^2}, \bar{z})| \\
 & \quad \times \left| \frac{\partial}{\partial z_1} W_t^{\lambda_1}(x_1, x_1 + \sqrt{\varepsilon^2 - |\bar{z}-\bar{x}|^2}) \right| \prod_{j=2}^n W_t^{\lambda_j}(x_j, z_j) \\
 & \quad + |(x_1 - \sqrt{\varepsilon^2 - |\bar{z}-\bar{x}|^2})^{2\lambda_1} f(x_1 - \sqrt{\varepsilon^2 - |\bar{z}-\bar{x}|^2}, \bar{z})| \\
 & \quad \times \left| \frac{\partial}{\partial z_1} W_t^{\lambda_1}(x_1, x_1 - \sqrt{\varepsilon^2 - |\bar{z}-\bar{x}|^2}) - \frac{\partial}{\partial z_1} W_t^{\lambda_1}(x_1, x_1 + \sqrt{\varepsilon^2 - |\bar{z}-\bar{x}|^2}) \right| \prod_{j=2}^n W_t^{\lambda_j}(x_j, z_j).
 \end{aligned}$$

By (22), mean value theorem leads to

$$\begin{aligned}
 & |(x_1 - \sqrt{\varepsilon^2 - |\bar{z}-\bar{x}|^2})^{2\lambda_1} f(x_1 - \sqrt{\varepsilon^2 - |\bar{z}-\bar{x}|^2}, \bar{z}) - (x_1 + \sqrt{\varepsilon^2 - |\bar{z}-\bar{x}|^2})^{2\lambda_1} f(x_1 + \sqrt{\varepsilon^2 - |\bar{z}-\bar{x}|^2}, \bar{z})| \\
 & \quad \times \left| \frac{\partial}{\partial z_1} W_t^{\lambda_1}(x_1, x_1 + \sqrt{\varepsilon^2 - |\bar{z}-\bar{x}|^2}) \right| \prod_{j=2}^n W_t^{\lambda_j}(x_j, z_j) \\
 (27) \quad & \leq C \frac{\varepsilon}{t^{\lambda_1+3/2}} \prod_{j=2}^n \frac{1}{t^{\lambda_j+1/2}}, \quad t \geq 1, \quad \bar{z} \in K^{n-1} \text{ and } |\bar{z}-\bar{x}| < \varepsilon.
 \end{aligned}$$

On the other hand, by (6) we get

$$\begin{aligned}
 \frac{\partial^2}{\partial z_1^2} W_t^{\lambda_1}(x_1, z_1) &= \frac{e^{-(x_1^2+z_1^2)/4t}}{(2t)^{\lambda_1+1/2}} \left\{ \left(\frac{x_1}{2t} \right)^2 \left[\left(\frac{x_1 z_1}{2t} \right)^{-\lambda_1-1/2} I_{\lambda_1+1/2} \left(\frac{x_1 z_1}{2t} \right) \right. \right. \\
 & \quad \left. \left. + \left(\frac{x_1 z_1}{2t} \right)^2 \left(\frac{x_1 z_1}{2t} \right)^{-\lambda_1-3/2} I_{\lambda_1+3/2} \left(\frac{x_1 z_1}{2t} \right) - \frac{z_1^2}{2t} \left(\frac{x_1 z_1}{2t} \right)^{-\lambda_1-1/2} I_{\lambda_1+1/2} \left(\frac{x_1 z_1}{2t} \right) \right] \right. \\
 & \quad \left. - \frac{1}{2t} \left(\frac{x_1 z_1}{2t} \right)^{-\lambda_1+1/2} I_{\lambda_1-1/2} \left(\frac{x_1 z_1}{2t} \right) \right. \\
 & \quad \left. - \frac{z_1}{2t} \left[\frac{x_1^2 z_1}{4t^2} \left(\frac{x_1 z_1}{2t} \right)^{-\lambda_1-1/2} I_{\lambda_1+1/2} \left(\frac{x_1 z_1}{2t} \right) - \frac{z_1}{2t} \left(\frac{x_1 z_1}{2t} \right)^{-\lambda_1+1/2} I_{\lambda_1-1/2} \left(\frac{x_1 z_1}{2t} \right) \right] \right\}, \quad t, z_1 > 0.
 \end{aligned}$$

Then, (4) implies that

$$\begin{aligned} \left| \frac{\partial^2}{\partial z_1^2} W_t^{\lambda_1}(x_1, z_1) \right| &\leq C \frac{e^{-(x_1^2 + z_1^2)/4t}}{t^{\lambda_1 + 1/2}} \\ &\times \left(\frac{x_1^2}{t^2} \left[1 + \left(\frac{x_1 z_1}{t} \right)^2 + \frac{z_1^2}{t} \right] + \frac{1}{t} + \frac{z_1}{t} \left(\frac{x_1^2 z_1}{t^2} + \frac{z_1}{t} \right) \right) \leq \frac{C}{t^{\lambda_1 + 3/2}}, \quad t \geq 1, \quad z_1 \in K. \end{aligned}$$

Hence, it obtains

$$\begin{aligned} & \left| (x_1 - \sqrt{\varepsilon^2 - |\bar{z} - \bar{x}|^2})^{2\lambda_1} f(x_1 - \sqrt{\varepsilon^2 - |\bar{z} - \bar{x}|^2}, \bar{z}) \right. \\ & \quad \times \left| \frac{\partial}{\partial z_1} W_t^{\lambda_1}(x_1, x_1 - \sqrt{\varepsilon^2 - |\bar{z} - \bar{x}|^2}) - \frac{\partial}{\partial z_1} W_t^{\lambda_1}(x_1, x_1 + \sqrt{\varepsilon^2 - |\bar{z} - \bar{x}|^2}) \right| \prod_{j=2}^n W_t^{\lambda_j}(x_j, z_j) \\ (28) \quad & \leq C \frac{\varepsilon}{t^{\lambda_1 + 3/2}} \prod_{j=2}^n \frac{1}{t^{\lambda_j + 1/2}}, \quad t \geq 1, \quad \bar{z} \in K^{n-1} \text{ and } |\bar{z} - \bar{x}| < \varepsilon. \end{aligned}$$

From (27) and (28) we deduce that

$$\begin{aligned} & \int_1^\infty |\phi(t)| \int_{\bar{z} \in K^{n-1}, |\bar{z} - \bar{x}| < \varepsilon} \left| H_2(x, x_1 - \sqrt{\varepsilon^2 - |\bar{z} - \bar{x}|^2}, \bar{z}, t) - H_2(x, x_1 + \sqrt{\varepsilon^2 - |\bar{z} - \bar{x}|^2}, \bar{z}, t) \right| \prod_{j=2}^n z_j^{2\lambda_j} d\bar{z} dt \\ (29) \quad & \leq C\varepsilon \int_1^\infty \int_{\bar{z} \in K^{n-1}, |\bar{z} - \bar{x}| < \varepsilon} \frac{1}{\sum_{j=1}^n (\lambda_j + 1/2) + 1} \prod_{j=2}^n z_j^{2\lambda_j} d\bar{z} dt \longrightarrow 0, \text{ as } \varepsilon \rightarrow 0^+. \end{aligned}$$

By (5) and (22) it has,

$$\begin{aligned} & \left| (x_1 - \sqrt{\varepsilon^2 - |\bar{z} - \bar{x}|^2})^{2\lambda_1} f(x_1 - \sqrt{\varepsilon^2 - |\bar{z} - \bar{x}|^2}, \bar{z}) - (x_1 + \sqrt{\varepsilon^2 - |\bar{z} - \bar{x}|^2})^{2\lambda_1} f(x_1 + \sqrt{\varepsilon^2 - |\bar{z} - \bar{x}|^2}, \bar{z}) \right| \\ & \quad \times \left| \frac{\partial}{\partial z_1} W_t^{\lambda_1}(x_1, x_1 + \sqrt{\varepsilon^2 - |\bar{z} - \bar{x}|^2}) \right| \prod_{j=2}^n W_t^{\lambda_j}(x_j, z_j) \\ & \leq C \sqrt{\varepsilon^2 - |\bar{z} - \bar{x}|^2} \frac{1}{t^{(n+1)/2}} \prod_{j=2}^n (x_j z_j)^{-\lambda_j} e^{-\frac{|\bar{x} - \bar{z}|^2}{4t} - \frac{\varepsilon^2 - |\bar{x} - \bar{z}|^2}{8t}} \\ & \leq C\varepsilon \frac{1}{t^{(n+1)/2}} e^{-\frac{\varepsilon^2}{8t}}, \quad 0 < t < 1, \quad \bar{z} \in K^{n-1} \text{ and } |\bar{z} - \bar{x}| < \varepsilon. \end{aligned}$$

Then [20, Lemma 1.1] allows us to get

$$\begin{aligned} & \int_0^1 |\phi(t)| \int_{|\bar{x} - \bar{z}| < \varepsilon} \left| (x_1 - \sqrt{\varepsilon^2 - |\bar{z} - \bar{x}|^2})^{2\lambda_1} f(x_1 - \sqrt{\varepsilon^2 - |\bar{z} - \bar{x}|^2}, \bar{z}) \right. \\ & \quad \left. - (x_1 + \sqrt{\varepsilon^2 - |\bar{z} - \bar{x}|^2})^{2\lambda_1} f(x_1 + \sqrt{\varepsilon^2 - |\bar{z} - \bar{x}|^2}, \bar{z}) \right| \left| \frac{\partial}{\partial z_1} W_t^{\lambda_1}(x_1, x_1 + \sqrt{\varepsilon^2 - |\bar{z} - \bar{x}|^2}) \right| \\ & \quad \times \prod_{j=2}^n W_t^{\lambda_j}(x_j, z_j) \prod_{j=2}^n z_j^{2\lambda_j} d\bar{z} dt \\ & \leq C\varepsilon \int_{|\bar{x} - \bar{z}| < \varepsilon} \int_0^1 \frac{e^{-\varepsilon^2/8t}}{t^{(n+1)/2}} dt d\bar{z} \\ & \leq C\varepsilon \longrightarrow 0, \quad \text{as } \varepsilon \rightarrow 0^+. \end{aligned}$$

Also, we write, for each $0 < t < 1$, $\bar{z} \in K^{n-1}$ and $|\bar{z} - \bar{x}| < \varepsilon$,

$$\begin{aligned}
& (x_1 - \sqrt{\varepsilon^2 - |\bar{z} - \bar{x}|^2})^{2\lambda_1} f(x_1 - \sqrt{\varepsilon^2 - |\bar{z} - \bar{x}|^2}, \bar{z}) \left(\frac{\partial}{\partial z_1} W_t^{\lambda_1}(x_1, x_1 - \sqrt{\varepsilon^2 - |\bar{z} - \bar{x}|^2}) \right. \\
& \quad \left. - \frac{\partial}{\partial z_1} W_t^{\lambda_1}(x_1, x_1 + \sqrt{\varepsilon^2 - |\bar{z} - \bar{x}|^2}) \right) \prod_{j=2}^n W_t^{\lambda_j}(x_j, z_j) \\
&= \left((x_1 - \sqrt{\varepsilon^2 - |\bar{z} - \bar{x}|^2})^{2\lambda_1} f(x_1 - \sqrt{\varepsilon^2 - |\bar{z} - \bar{x}|^2}, \bar{z}) - x_1^{2\lambda_1} f(x) \right) \left(\frac{\partial}{\partial z_1} W_t^{\lambda_1}(x_1, x_1 - \sqrt{\varepsilon^2 - |\bar{z} - \bar{x}|^2}) \right. \\
& \quad \left. - \frac{\partial}{\partial z_1} W_t^{\lambda_1}(x_1, x_1 + \sqrt{\varepsilon^2 - |\bar{z} - \bar{x}|^2}) \right) \prod_{j=2}^n W_t^{\lambda_j}(x_j, z_j) + x_1^{2\lambda_1} f(x) \left(\frac{\partial}{\partial z_1} W_t^{\lambda_1}(x_1, x_1 - \sqrt{\varepsilon^2 - |\bar{z} - \bar{x}|^2}) \right. \\
& \quad \left. - \frac{\partial}{\partial z_1} W_t^{\lambda_1}(x_1, x_1 + \sqrt{\varepsilon^2 - |\bar{z} - \bar{x}|^2}) \right) \prod_{j=2}^n W_t^{\lambda_j}(x_j, z_j).
\end{aligned}$$

By (22) we have

$$\left| \frac{\partial}{\partial z_1} W_t^{\lambda_1}(x_1, z_1) \right| \leq C \frac{e^{-(x_1 - z_1)^2/8t}}{t}, \quad 0 < t < 1, \quad z_1 \in K.$$

By proceeding as above we obtain,

$$\begin{aligned}
& \int_0^1 |\phi(t)| \int_{\bar{z} \in K^{n-1}, |\bar{x} - \bar{z}| < \varepsilon} \left| (x_1 - \sqrt{\varepsilon^2 - |\bar{z} - \bar{x}|^2})^{2\lambda_1} f(x_1 - \sqrt{\varepsilon^2 - |\bar{z} - \bar{x}|^2}, \bar{z}) - x_1^{2\lambda_1} f(x) \right| \\
& \quad \times \left| \frac{\partial}{\partial z_1} W_t^{\lambda_1}(x_1, x_1 - \sqrt{\varepsilon^2 - |\bar{z} - \bar{x}|^2}) - \frac{\partial}{\partial z_1} W_t^{\lambda_1}(x_1, x_1 + \sqrt{\varepsilon^2 - |\bar{z} - \bar{x}|^2}) \right| \prod_{j=2}^n W_t^{\lambda_j}(x_j, z_j) \prod_{j=2}^n z_j^{2\lambda_j} d\bar{z} \\
& \leq C\varepsilon \int_{\bar{z} \in K^{n-1}, |\bar{x} - \bar{z}| < \varepsilon} \int_0^1 \frac{e^{-\varepsilon^2/8t}}{t^{(n+1)/2}} dt d\bar{z} \\
& \leq C\varepsilon \longrightarrow 0, \quad \text{as } \varepsilon \rightarrow 0^+.
\end{aligned}$$

From the above estimates we conclude that

$$\begin{aligned}
& \int_0^\infty \phi(t) \int_{(0, \infty)^n} \Delta_{\lambda_1, z_1} f(z) \left(\prod_{j=1}^n W_t^{\lambda_j}(x_j, z_j) - \frac{\chi_{(1, \infty)}(t) t^{-\sum_{j=1}^n (\lambda_j + 1/2)}}{\prod_{j=1}^n 2^{2\lambda_j} \Gamma(\lambda_j + 1/2)} \right) \prod_{j=1}^n z_j^{2\lambda_j} dz dt \\
&= - \lim_{\varepsilon \rightarrow 0^+} \left[\int_0^\infty \phi(t) \int_{|x-z| > \varepsilon} f(z) \frac{\partial}{\partial t} \left(W_t^{\lambda_1}(x_1, z_1) \right) \prod_{j=2}^n W_t^{\lambda_j}(x_j, z_j) \prod_{j=1}^n z_j^{2\lambda_j} dz dt \right. \\
& \quad \left. + f(x) x_1^{2\lambda_1} \int_0^1 \phi(t) \int_{|\bar{x} - \bar{z}| < \varepsilon} \int_{x_1 - \sqrt{\varepsilon^2 - |\bar{x} - \bar{z}|^2}}^{x_1 + \sqrt{\varepsilon^2 - |\bar{x} - \bar{z}|^2}} \frac{\partial^2}{\partial z_1^2} W_t^{\lambda_1}(x_1, z_1) \prod_{j=2}^n W_t^{\lambda_j}(x_j, z_j) dz_1 \prod_{j=2}^n z_j^{2\lambda_j} d\bar{z} dt \right] \\
&= - \lim_{\varepsilon \rightarrow 0^+} \left[\int_0^\infty \phi(t) \int_{|x-z| > \varepsilon} f(z) \frac{\partial}{\partial t} \left(W_t^{\lambda_1}(x_1, z_1) \right) \prod_{j=2}^n W_t^{\lambda_j}(x_j, z_j) \prod_{j=1}^n z_j^{2\lambda_j} dz dt \right. \\
& \quad \left. + f(x) x_1^{2\lambda_1} \int_0^1 \phi(t) \int_{|x-z| < \varepsilon} \frac{\partial^2}{\partial z_1^2} W_t^{\lambda_1}(x_1, z_1) \prod_{j=2}^n W_t^{\lambda_j}(x_j, z_j) \prod_{j=2}^n z_j^{2\lambda_j} dz dt \right].
\end{aligned}$$

We now denote by $\mathbb{W}_t(u, v)$ the classical heat kernel, that is,

$$\mathbb{W}_t(u, v) = \frac{1}{2\sqrt{\pi t}} e^{-|u-v|^2/4t}, \quad u, v \in \mathbb{R}, \quad t > 0.$$

According to (6) we get

$$\begin{aligned}
\frac{\partial^2}{\partial z_1^2} W_t^{\lambda_1}(x_1, z_1) &= \frac{\partial^2}{\partial z_1^2} \left[\frac{\sqrt{2\pi}}{(2t)^{\lambda_1}} \left(\frac{x_1 z_1}{2t} \right)^{-\lambda_1+1/2} I_{\lambda_1-1/2} \left(\frac{x_1 z_1}{2t} \right) e^{-x_1 z_1/2t} \mathbb{W}_t(x_1, z_1) \right] \\
&= \frac{\sqrt{2\pi}}{(2t)^{\lambda_1}} \left\{ \frac{\partial^2}{\partial z_1^2} \left[\left(\frac{x_1 z_1}{2t} \right)^{-\lambda_1+1/2} I_{\lambda_1-1/2} \left(\frac{x_1 z_1}{2t} \right) e^{-x_1 z_1/2t} \right] \mathbb{W}_t(x_1, z_1) \right. \\
&\quad + 2 \frac{\partial}{\partial z_1} \left[\left(\frac{x_1 z_1}{2t} \right)^{-\lambda_1+1/2} I_{\lambda_1-1/2} \left(\frac{x_1 z_1}{2t} \right) e^{-x_1 z_1/2t} \right] \frac{\partial}{\partial z_1} \mathbb{W}_t(x_1, z_1) \\
&\quad \left. + \left(\frac{x_1 z_1}{2t} \right)^{-\lambda_1+1/2} I_{\lambda_1-1/2} \left(\frac{x_1 z_1}{2t} \right) e^{-x_1 z_1/2t} \frac{\partial^2}{\partial z_1^2} \mathbb{W}_t(x_1, z_1) \right\} \\
&= \frac{\sqrt{2\pi}}{(2t)^{\lambda_1}} e^{-x_1 z_1/2t} \left(\frac{x_1 z_1}{2t} \right)^{-\lambda_1} \left\{ \left[\left(\frac{x_1 z_1}{2t} \right)^{1/2} I_{\lambda_1+3/2} \left(\frac{x_1 z_1}{2t} \right) \right. \right. \\
&\quad + \left(\frac{x_1 z_1}{2t} \right)^{1/2} I_{\lambda_1+1/2} \left(\frac{x_1 z_1}{2t} \right) \frac{2t}{x_1 z_1} - 2 \left(\frac{x_1 z_1}{2t} \right)^{1/2} I_{\lambda_1+1/2} \left(\frac{x_1 z_1}{2t} \right) \\
&\quad \left. + \left(\frac{x_1 z_1}{2t} \right)^{1/2} I_{\lambda_1-1/2} \left(\frac{x_1 z_1}{2t} \right) \right] \mathbb{W}_t(x_1, z_1) \left(\frac{x_1}{2t} \right)^2 \\
&\quad + \frac{x_1}{t} \left[\left(\frac{x_1 z_1}{2t} \right)^{1/2} I_{\lambda_1+1/2} \left(\frac{x_1 z_1}{2t} \right) - \left(\frac{x_1 z_1}{2t} \right)^{1/2} I_{\lambda_1-1/2} \left(\frac{x_1 z_1}{2t} \right) \right] \frac{\partial}{\partial z_1} \mathbb{W}_t(x_1, z_1) \\
&\quad \left. + \left[\left(\frac{x_1 z_1}{2t} \right)^{1/2} I_{\lambda_1-1/2} \left(\frac{x_1 z_1}{2t} \right) \right] \frac{\partial^2}{\partial z_1^2} \mathbb{W}_t(x_1, z_1) \right\}, \quad t, z_1 \in (0, \infty).
\end{aligned}$$

By using (5) we obtain for every $0 < t < 1$, and $z_1 \in K$,

$$\begin{aligned}
&\bullet \left| \frac{\sqrt{2\pi}}{(2t)^{\lambda_1}} \left(\frac{x_1 z_1}{2t} \right)^{-\lambda_1+1/2} I_{\lambda_1-1/2} \left(\frac{x_1 z_1}{2t} \right) e^{-x_1 z_1/2t} \frac{\partial^2}{\partial z_1^2} \mathbb{W}_t(x_1, z_1) - (x_1 z_1)^{-\lambda_1} \frac{\partial^2}{\partial z_1^2} \mathbb{W}_t(x_1, z_1) \right| \\
(30) \quad &\leq C t \left| \frac{\partial^2}{\partial z_1^2} \mathbb{W}_t(x_1, z_1) \right|,
\end{aligned}$$

$$\begin{aligned}
&\bullet \left| \frac{\sqrt{2\pi}}{(2t)^{\lambda_1}} \left(\frac{x_1 z_1}{2t} \right)^{-\lambda_1} \frac{x_1}{2t} \frac{\partial}{\partial z_1} \mathbb{W}_t(x_1, z_1) e^{-x_1 z_1/2t} \left[\left(\frac{x_1 z_1}{2t} \right)^{1/2} I_{\lambda_1+1/2} \left(\frac{x_1 z_1}{2t} \right) \right. \right. \\
(31) \quad &\quad \left. \left. - \left(\frac{x_1 z_1}{2t} \right)^{1/2} I_{\lambda_1-1/2} \left(\frac{x_1 z_1}{2t} \right) \right] \right| \leq C \left| \frac{\partial}{\partial z_1} \mathbb{W}_t(x_1, z_1) \right|;
\end{aligned}$$

and

$$\begin{aligned}
&\bullet \left| \frac{\sqrt{2\pi}}{(2t)^{\lambda_1}} e^{-x_1 z_1/2t} \left(\frac{x_1 z_1}{2t} \right)^{-\lambda_1} \mathbb{W}_t(x_1, z_1) \left(\frac{x_1}{2t} \right)^2 \left(\left(\frac{x_1 z_1}{2t} \right)^{1/2} I_{\lambda_1+3/2} \left(\frac{x_1 z_1}{2t} \right) \right. \right. \\
&\quad + \left(\frac{x_1 z_1}{2t} \right)^{1/2} I_{\lambda_1+1/2} \left(\frac{x_1 z_1}{2t} \right) \frac{2t}{x_1 z_1} - 2 \left(\frac{x_1 z_1}{2t} \right)^{1/2} I_{\lambda_1+1/2} \left(\frac{x_1 z_1}{2t} \right) \\
&\quad \left. + \left(\frac{x_1 z_1}{2t} \right)^{1/2} I_{\lambda_1-1/2} \left(\frac{x_1 z_1}{2t} \right) \right) \right| \\
(32) \quad &\leq C \mathbb{W}_t(x_1, z_1).
\end{aligned}$$

Hence, we get for every $0 < t < 1$, and $z_1 \in K$,

$$\left| \frac{\partial^2}{\partial z_1^2} W_t^{\lambda_1}(x_1, z_1) - (x_1 z_1)^{-\lambda_1} \frac{\partial^2}{\partial z_1^2} \mathbb{W}_t(x_1, z_1) \right| \leq C \frac{e^{-(x_1 - z_1)^2/8t}}{t}.$$

Then, by involving again (5) and [20, Lemma 1.1] we have that

$$\begin{aligned}
& \int_0^1 |\phi(t)| \int_{|x-z|<\varepsilon} \left| \frac{\partial^2}{\partial z_1^2} W_t^{\lambda_1}(x_1, z_1) - (x_1 z_1)^{-\lambda_1} \frac{\partial^2}{\partial z_1^2} \mathbb{W}_t(x_1, z_1) \right| \prod_{j=2}^n W_t^{\lambda_j}(x_j, z_j) \prod_{j=2}^n z_j^{2\lambda_j} dz dt \\
& \leq C \int_{|x-z|<\varepsilon} \int_0^1 \frac{e^{-|x-z|^2/8t}}{t^{(n+1)/2}} \prod_{j=2}^n z_j^{2\lambda_j} dt dz \\
& \leq C \int_{|x-z|<\varepsilon} \frac{dz}{|x-z|^{n-1}} \leq C\varepsilon \longrightarrow 0, \text{ as } \varepsilon \rightarrow 0^+.
\end{aligned}$$

Also, (5) leads to, for every $0 < t < 1$, and $z \in K^n$,

$$\begin{aligned}
& \left| \prod_{j=1}^{i-1} (x_j z_j)^{-\lambda_j} \frac{\partial^2}{\partial z_1^2} \mathbb{W}_t(x_1, z_1) \prod_{j=2}^{i-1} \mathbb{W}_t(x_j, z_j) \left(W_t^{\lambda_i}(x_i, z_i) - (x_i z_i)^{-\lambda_i} \mathbb{W}_t(x_i, z_i) \right) \right. \\
& \quad \left. \times \prod_{j=i+1}^n W_t^{\lambda_j}(x_j, z_j) \right| \leq C \frac{e^{-|x-z|^2/8t}}{t^{n/2}}, \quad i = 2, \dots, n.
\end{aligned}$$

Then, we have that for every $i = 2, \dots, n$,

$$\begin{aligned}
& \int_0^1 |\phi(t)| \int_{|x-z|<\varepsilon} \left| \prod_{j=1}^{i-1} (x_j z_j)^{-\lambda_j} \frac{\partial^2}{\partial z_1^2} \mathbb{W}_t(x_1, z_1) \prod_{j=2}^{i-1} \mathbb{W}_t(x_j, z_j) \left(W_t^{\lambda_i}(x_i, z_i) - (x_i z_i)^{-\lambda_i} \mathbb{W}_t(x_i, z_i) \right) \right. \\
& \quad \left. \times \prod_{j=i+1}^n W_t^{\lambda_j}(x_j, z_j) \right| \prod_{j=2}^n z_j^{2\lambda_j} dz dt \\
& \leq C \int_{|x-z|<\varepsilon} \int_0^1 \frac{e^{-|x-z|^2/8t}}{t^{n/2}} dt dz \\
& \leq C \int_{|x-z|<\varepsilon} \frac{dz}{|x-z|^{n-2}} \longrightarrow 0, \text{ as } \varepsilon \rightarrow 0^+,
\end{aligned}$$

provided that $n > 2$. When $n = 2$ we proceed in a similar way.

Hence, we conclude that

$$\begin{aligned}
& \int_0^\infty \phi(t) \int_{(0,\infty)^n} \Delta_{\lambda_1, z_1} f(z) \left(\prod_{j=1}^n W_t^{\lambda_j}(x_j, z_j) - \frac{\chi_{(1,\infty)}(t) t^{-\sum_{j=1}^n (\lambda_j+1/2)}}{\prod_{j=1}^n 2^{2\lambda_j} \Gamma(\lambda_j+1/2)} \right) \prod_{j=1}^n z_j^{2\lambda_j} dz_j dt \\
& = - \lim_{\varepsilon \rightarrow 0^+} \left[\int_0^\infty \phi(t) \int_{|x-z|>\varepsilon} f(z) \frac{\partial}{\partial t} \left(W_t^{\lambda_1}(x_1, z_1) \right) \prod_{j=2}^n W_t^{\lambda_j}(x_j, z_j) \prod_{j=1}^n z_j^{2\lambda_j} dz dt \right. \\
& \quad \left. + f(x) x_1^{2\lambda_1} \int_0^1 \phi(t) \int_{|x-z|<\varepsilon} \prod_{j=1}^n (x_j z_j)^{-\lambda_j} \frac{\partial^2}{\partial z_1^2} \mathbb{W}_t(x_1, z_1) \prod_{j=2}^n \mathbb{W}_t(x_j, z_j) \prod_{j=2}^n z_j^{2\lambda_j} dz dt \right].
\end{aligned}$$

On the other hand, the mean value theorem allows us to write, for every $\alpha \in \mathbb{R}$, and $j = 1, \dots, n$,

$$\begin{aligned}
& |z_j^\alpha - x_j^\alpha| \left| \frac{\partial^2}{\partial z_1^2} \mathbb{W}_t(x_1, z_1) \prod_{j=2}^n \mathbb{W}_t(x_j, z_j) \right| \leq C |z_j - x_j| \frac{e^{-|x-z|^2/8t}}{t^{n/2+1}} \\
& \leq C \frac{e^{-|x-z|^2/16t}}{t^{(n+1)/2}}, \quad t > 0 \text{ and } z \in K^n.
\end{aligned}$$

Then, by proceeding as above we obtain

$$\begin{aligned}
& \int_0^\infty \phi(t) \int_{(0,\infty)^n} \Delta_{\lambda_1, z_1} f(z) \left(\prod_{j=1}^n W_t^{\lambda_j}(x_j, z_j) - \frac{\chi_{(1,\infty)}(t)t^{-\sum_{j=1}^n (\lambda_j+1/2)}}{\prod_{j=1}^n 2^{2\lambda_j} \Gamma(\lambda_j + 1/2)} \right) \prod_{j=1}^n z_j^{2\lambda_j} dz_j dt \\
&= - \lim_{\varepsilon \rightarrow 0^+} \left[\int_0^\infty \phi(t) \int_{|x-z|>\varepsilon} f(z) \frac{\partial}{\partial t} \left(W_t^{\lambda_1}(x_1, z_1) \right) \prod_{j=2}^n W_t^{\lambda_j}(x_j, z_j) \prod_{j=1}^n z_j^{2\lambda_j} dz dt \right. \\
&\quad \left. + f(x) \int_0^1 \phi(t) \int_{|x-z|<\varepsilon} \frac{\partial^2}{\partial z_1^2} \frac{e^{-|x-z|^2/4t}}{(2\sqrt{\pi t})^n} dz dt \right] \\
&= - \lim_{\varepsilon \rightarrow 0^+} \left[\int_0^\infty \phi(t) \int_{|x-z|>\varepsilon} f(z) \frac{\partial}{\partial t} \left(W_t^{\lambda_1}(x_1, z_1) \right) \prod_{j=2}^n W_t^{\lambda_j}(x_j, z_j) \prod_{j=1}^n z_j^{2\lambda_j} dz dt \right. \\
(33) \quad &\left. + f(x) \int_0^1 \phi(t) \int_{|y|<\varepsilon} \frac{\partial^2}{\partial y_1^2} \frac{e^{-|y|^2/4t}}{(2\sqrt{\pi t})^n} dy dt \right].
\end{aligned}$$

Also, we have that, for every $i = 2, \dots, n$,

$$\begin{aligned}
& \int_0^\infty \phi(t) \int_{(0,\infty)^n} \Delta_{\lambda_i, z_i} f(z) \left(\prod_{j=1}^n W_t^{\lambda_j}(x_j, z_j) - \frac{\chi_{(1,\infty)}(t)t^{-\sum_{j=1}^n (\lambda_j+1/2)}}{\prod_{j=1}^n 2^{2\lambda_j} \Gamma(\lambda_j + 1/2)} \right) \prod_{j=1}^n z_j^{2\lambda_j} dz_j dt \\
(34) \quad &= - \lim_{\varepsilon \rightarrow 0^+} \left[\int_0^\infty \phi(t) \int_{|x-z|>\varepsilon} f(z) \frac{\partial}{\partial t} \left(W_t^{\lambda_i}(x_i, z_i) \right) \prod_{j=1, j \neq i}^n W_t^{\lambda_j}(x_j, z_j) \prod_{j=1}^n z_j^{2\lambda_j} dz dt \right. \\
&\quad \left. + f(x) \int_0^1 \phi(t) \int_{|y|<\varepsilon} \frac{\partial^2}{\partial y_i^2} \frac{e^{-|y|^2/4t}}{(2\sqrt{\pi t})^n} dy dt \right].
\end{aligned}$$

Then, by (33) and (34), it follows that

$$\begin{aligned}
(35) \quad T_{\lambda_1, \dots, \lambda_n}^m f(x) &= - \lim_{\varepsilon \rightarrow 0^+} \left[\int_0^\infty \phi(t) \int_{|x-z|>\varepsilon} f(z) \frac{\partial}{\partial t} \left(\prod_{j=1}^n W_t^{\lambda_j}(x_j, z_j) \right) \prod_{j=1}^n z_j^{2\lambda_j} dz dt \right. \\
&\quad \left. + n f(x) \int_0^1 \phi(t) \int_{|y|<\varepsilon} \frac{\partial^2}{\partial y_1^2} \frac{e^{-|y|^2/4t}}{(2\sqrt{\pi t})^n} dy dt \right].
\end{aligned}$$

We define

$$\alpha(\varepsilon) = \int_0^1 \phi(t) \int_{|y|<\varepsilon} \frac{\partial^2}{\partial y_1^2} \frac{e^{-|y|^2/4t}}{(2\sqrt{\pi t})^n} dy dt, \quad \varepsilon > 0.$$

We can write

$$\begin{aligned}
\alpha(\varepsilon) &= \int_0^1 \phi(t) \int_{|\bar{y}|<\varepsilon} \int_{-\sqrt{\varepsilon^2-|\bar{y}|^2}}^{\sqrt{\varepsilon^2-|\bar{y}|^2}} \frac{\partial^2}{\partial y_1^2} \frac{e^{-|y|^2/4t}}{(2\sqrt{\pi t})^n} dy_1 d\bar{y} dt \\
&= \int_0^1 \phi(t) \int_{|\bar{y}|<\varepsilon} \left[\frac{\partial}{\partial y_1} \left(\frac{e^{-(y_1^2+|\bar{y}|^2)/4t}}{(2\sqrt{\pi t})^n} \right) \Big|_{y_1=\sqrt{\varepsilon^2-|\bar{y}|^2}} - \frac{\partial}{\partial y_1} \left(\frac{e^{-(y_1^2+|\bar{y}|^2)/4t}}{(2\sqrt{\pi t})^n} \right) \Big|_{y_1=-\sqrt{\varepsilon^2-|\bar{y}|^2}} \right] d\bar{y} dt, \quad \varepsilon > 0.
\end{aligned}$$

Hence, [20, Lemma 1.1] leads to

$$|\alpha(\varepsilon)| \leq C \int_0^1 \int_{|\bar{y}|<\varepsilon} \frac{e^{-\varepsilon^2/8t}}{t^{(n+1)/2}} d\bar{y} dt \leq C \int_{|\bar{y}|<\varepsilon} \int_0^1 \frac{e^{-\varepsilon^2/8t}}{t^{(n+1)/2}} dt d\bar{y} \leq \frac{1}{\varepsilon^{n-1}} \int_{|\bar{y}|<\varepsilon} d\bar{y} \leq C, \quad \varepsilon > 0.$$

Suppose now that there exists $\phi(0^+) = \lim_{t \rightarrow 0^+} \phi(t)$. Then, we have that

$$(36) \quad \lim_{\varepsilon \rightarrow 0^+} \int_0^1 \phi(t) \int_{|y|<\varepsilon} \frac{\partial^2}{\partial y_1^2} \frac{e^{-|y|^2/4t}}{(2\sqrt{\pi t})^n} dy dt = -M\phi(0^+),$$

for a certain $M > 0$. Indeed, by making changes of variables we obtain

$$\begin{aligned}
\int_0^1 \phi(t) \int_{|y|<\varepsilon} \frac{\partial^2}{\partial y_1^2} \frac{e^{-|y|^2/4t}}{(2\sqrt{\pi t})^n} dy dt &= \int_0^{1/\varepsilon^2} \phi(s\varepsilon^2) \int_{|z|<1} \frac{\partial^2}{\partial z_1^2} \frac{e^{-|z|^2/4s}}{(2\sqrt{\pi s})^n} dz ds \\
&= \int_0^{1/\varepsilon^2} \phi(s\varepsilon^2) \int_{|\bar{z}|<1} \frac{\partial}{\partial z_1} \frac{e^{-|z|^2/4s}}{(2\sqrt{\pi s})^n} \Bigg]_{z_1=-\sqrt{1-|\bar{z}|^2}}^{z_1=\sqrt{1-|\bar{z}|^2}} d\bar{z} ds \\
&= - \int_0^{1/\varepsilon^2} \phi(s\varepsilon^2) \int_{|\bar{z}|<1} \frac{z_1}{2s} \frac{e^{-|z|^2/4s}}{(2\sqrt{\pi s})^n} \Bigg]_{z_1=-\sqrt{1-|\bar{z}|^2}}^{z_1=\sqrt{1-|\bar{z}|^2}} d\bar{z} ds \\
&= - \int_0^{1/\varepsilon^2} \phi(s\varepsilon^2) \int_{|\bar{z}|<1} \frac{\sqrt{1-|\bar{z}|^2}}{s} \frac{e^{-1/4s}}{(2\sqrt{\pi s})^n} d\bar{z} ds \\
&= -M \int_0^{1/\varepsilon^2} \phi(s\varepsilon^2) \frac{e^{-1/4s}}{s^{n/2+1}} ds, \quad \varepsilon > 0,
\end{aligned}$$

where

$$M = \frac{1}{(2\sqrt{\pi})^n} \int_{|\bar{z}|<1} \sqrt{1-|\bar{z}|^2} d\bar{z}.$$

Moreover, denoting by $\chi_{[0,1]}$ the characteristic function of $[0, 1]$, we have that

$$\int_0^{1/\varepsilon^2} \phi(s\varepsilon^2) \frac{e^{-1/4s}}{s^{n/2+1}} ds = \int_0^\infty \phi(s\varepsilon^2) \chi_{[0,1]}(s\varepsilon^2) \frac{e^{-1/4s}}{s^{n/2+1}} ds, \quad \varepsilon > 0.$$

Then, by using the dominated convergence theorem, it follows that

$$\lim_{\varepsilon \rightarrow 0^+} \int_0^{1/\varepsilon^2} \phi(s\varepsilon^2) \frac{e^{-1/4s}}{s^{n/2+1}} ds = \phi(0^+) \int_0^\infty \frac{e^{-1/4s}}{s^{n/2+1}} ds.$$

Hence, there exists $M > 0$ for which (36) holds.

Let $\alpha > -1/2$. According to (6) it follows that

$$\begin{aligned}
(37) \quad \frac{\partial}{\partial t} W_t^\alpha(u, v) &= \frac{e^{-(u^2+v^2)/4t}}{2^{\alpha+1/2}} \left\{ -\frac{\alpha+1/2}{t^{\alpha+3/2}} \left(\frac{uv}{2t}\right)^{-\alpha+1/2} I_{\alpha-1/2}\left(\frac{uv}{2t}\right) \right. \\
&\quad \left. - \frac{uv}{2t^{\alpha+5/2}} \left(\frac{uv}{2t}\right)^{-\alpha+1/2} I_{\alpha+1/2}\left(\frac{uv}{2t}\right) + \frac{u^2+v^2}{4t^{\alpha+5/2}} \left(\frac{uv}{2t}\right)^{-\alpha+1/2} I_{\alpha-1/2}\left(\frac{uv}{2t}\right) \right\}, \quad u, v, t \in (0, \infty).
\end{aligned}$$

Then, by (4) we obtains

$$(38) \quad \left| \frac{\partial}{\partial t} W_t^\alpha(u, v) \right| \leq C \frac{1}{t^{\alpha+3/2}}, \quad \frac{uv}{t} \leq 1, \quad u, v, t \in (0, \infty),$$

and, by (5) we have

$$\begin{aligned}
(39) \quad \left| \frac{\partial}{\partial t} W_t^\alpha(u, v) \right| &\leq C \frac{e^{-(u-v)^2/4t}}{t^{3/2}} (uv)^{-\alpha} \left(\frac{|u-v|^2}{t} + 1 + \frac{u^2+v^2}{uv} \right) \\
&\leq C \frac{e^{-(u-v)^2/4t}}{t^{3/2}} (uv)^{-\alpha} \left(1 + \frac{u^2+v^2}{uv} \right), \quad \frac{uv}{t} \geq 1, \quad u, v, t \in (0, \infty).
\end{aligned}$$

For every $t > 0$ and $z \in (0, \infty)^n$,

$$\begin{aligned}
\frac{\partial}{\partial t} \prod_{j=1}^n W_t^{\lambda_j}(x_j, z_j) &= \sum_{i=1}^n \prod_{\substack{j=1 \\ j \neq i}}^n W_t^{\lambda_j}(x_j, z_j) \frac{\partial}{\partial t} W_t^{\lambda_i}(x_i, z_i) \\
&= \sum_{i=1}^n \prod_{\substack{j=1 \\ j \neq i}}^n W_t^{\lambda_j}(x_j, z_j) \frac{e^{-(x_i^2+z_i^2)/4t}}{2^{\lambda_i+1/2}} \left\{ -\frac{\lambda_i+1/2}{t^{\lambda_i+3/2}} \left(\frac{x_i z_i}{2t}\right)^{-\lambda_i+1/2} I_{\lambda_i-1/2}\left(\frac{x_i z_i}{2t}\right) \right. \\
&\quad \left. - \frac{x_i^2 z_i^2}{4t^{\lambda_i+7/2}} \left(\frac{x_i z_i}{2t}\right)^{-\lambda_i-1/2} I_{\lambda_i+1/2}\left(\frac{x_i z_i}{2t}\right) + \frac{x_i^2+z_i^2}{4t^{\lambda_i+5/2}} \left(\frac{x_i z_i}{2t}\right)^{-\lambda_i+1/2} I_{\lambda_i-1/2}\left(\frac{x_i z_i}{2t}\right) \right\}.
\end{aligned}$$

Hence, by using (4), (5), (38), and 39, we get, for every $z \in K^n$,

$$\begin{aligned} \left| \frac{\partial}{\partial t} \prod_{j=1}^n W_t^{\lambda_j}(x_j, z_j) \right| &\leq \sum_{i=1}^n \left| \frac{\partial}{\partial t} W_t^{\lambda_i}(x_i, z_i) \right| \prod_{j=1, j \neq i}^n W_t^{\lambda_j}(x_j, z_j) \\ &\leq C \begin{cases} t^{-\sum_{j=1}^n (\lambda_j + 1/2) + 1} & , \quad t > 1 \\ e^{-|x-z|^2/8t} t^{-(n+2)/2} & , \quad 0 < t \leq 1. \end{cases} \end{aligned}$$

Hence, since $f \in C_c^\infty((0, \infty)^n)$, it follows that, for every $\varepsilon > 0$

$$\int_0^\infty |\phi(t)| \int_{(0, \infty)^n, |x-z| > \varepsilon} |f(z)| \left| \frac{\partial}{\partial t} \left(\prod_{j=1}^n W_t^{\lambda_j}(x_j, z_j) \right) \right| \prod_{j=1}^n z_j^{2\lambda_j} dz dt < \infty.$$

The we can interchange the order of integration on the integrals in (35) and by using (36) we conclude that

$$T_{\lambda_1, \dots, \lambda_n}^m f(x) = - \lim_{\varepsilon \rightarrow 0^+} \int_{(0, \infty)^n, |z-x| > \varepsilon} f(z) \int_0^\infty \phi(t) \frac{\partial}{\partial t} \prod_{j=1}^n W_t^{\lambda_j}(x_j, z_j) dt \prod_{j=1}^n z_j^{2\lambda_j} dz + C \phi(0^+) f(x),$$

for a certain $C > 0$. □

We now prove the L^p -boundedness properties of the maximal operator associated with the principal value integral that appears in Proposition 2.1.

Proposition 2.2. *Let $\lambda_j > -1/2$, $j = 1, \dots, n$. The maximal operator $T_{\lambda_1, \dots, \lambda_n}^{m,*}$ defined by*

$$T_{\lambda_1, \dots, \lambda_n}^{m,*}(f)(x) = \sup_{\varepsilon > 0} \left| \int_{(0, \infty)^n, |x-y| > \varepsilon} f(y) K_{\lambda_1, \dots, \lambda_n}^\phi(x, y) \prod_{j=1}^n y_j^{2\lambda_j} dy \right|, \quad x \in (0, \infty)^n,$$

where

$$K_{\lambda_1, \dots, \lambda_n}^\phi(x, y) = \int_0^\infty \phi(t) \frac{\partial}{\partial t} \prod_{j=1}^n W_t^{\lambda_j}(x_j, y_j) dt, \quad x, y \in (0, \infty)^n,$$

is bounded from $L^p((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$ into itself, for every $1 < p < \infty$, and from $L^1((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$ into $L^{1,\infty}((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$.

Proof. In order to established the L^p -boundedness properties for the maximal operator $T_{\lambda_1, \dots, \lambda_n}^{m,*}$ we consider the operator

$$T_{loc, \lambda_1, \dots, \lambda_n}^{m,*}(f)(x) = \sup_{\varepsilon > 0} \left| \int_{L(x), |x-y| > \varepsilon} f(y) \prod_{j=1}^n (x_j y_j)^{-\lambda_j} H^\phi(x, y) \prod_{j=1}^n y_j^{2\lambda_j} dy \right|, \quad x \in (0, \infty)^n,$$

where, for every $x = (x_1, \dots, x_n) \in (0, \infty)^n$,

$$L(x) = \{y = (y_1, \dots, y_n) \in (0, \infty)^n : x_j/2 < y_j < 2x_j, \quad j = 1, \dots, n\},$$

and

$$H^\phi(x, y) = \int_0^\infty \phi(t) \frac{\partial}{\partial t} \left[\frac{e^{-|x-y|^2/4t}}{(4\pi t)^{n/2}} \right] dt, \quad x, y \in (0, \infty)^n.$$

We have that

$$\begin{aligned}
T_{\lambda_1, \dots, \lambda_n}^{m,*}(f)(x) &\leq \left| T_{\lambda_1, \dots, \lambda_n}^{m,*}(f)(x) - T_{loc, \lambda_1, \dots, \lambda_n}^{m,*}(f)(x) \right| + T_{loc, \lambda_1, \dots, \lambda_n}^{m,*}(f)(x) \\
&\leq \sup_{\varepsilon > 0} \left| \int_{(0, \infty)^n \setminus L(x), |x-y| > \varepsilon} f(y) K_{\lambda_1, \dots, \lambda_n}^\phi(x, y) \prod_{j=1}^n y_j^{2\lambda_j} dy \right| \\
&\quad + \sup_{\varepsilon > 0} \left| \int_{L(x), |x-y| > \varepsilon} f(y) \left(K_{\lambda_1, \dots, \lambda_n}^\phi(x, y) - \prod_{j=1}^n (x_j y_j)^{-\lambda_j} H^\phi(x, y) \right) \prod_{j=1}^n y_j^{2\lambda_j} dy \right| \\
&\quad + T_{loc, \lambda_1, \dots, \lambda_n}^{m,*}(f)(x) \\
&\leq \int_{(0, \infty)^n \setminus L(x)} |f(y)| |K_{\lambda_1, \dots, \lambda_n}^\phi(x, y)| \prod_{j=1}^n y_j^{2\lambda_j} dy \\
&\quad + \int_{L(x)} |f(y)| \left| K_{\lambda_1, \dots, \lambda_n}^\phi(x, y) - \prod_{j=1}^n (x_j y_j)^{-\lambda_j} H^\phi(x, y) \right| \prod_{j=1}^n y_j^{2\lambda_j} dy \\
&\quad + T_{loc, \lambda_1, \dots, \lambda_n}^{m,*}(f)(x) \\
(40) \quad &= \mathcal{G}_{\lambda_1, \dots, \lambda_n}^\phi(|f|)(x) + \mathcal{L}_{\lambda_1, \dots, \lambda_n}(|f|)(x) + T_{loc, \lambda_1, \dots, \lambda_n}^{m,*}(f)(x), \quad x \in (0, \infty)^n.
\end{aligned}$$

We are going to show the L^p -boundedness properties for the operator $\mathcal{G}_{\lambda_1, \dots, \lambda_n}^\phi$, $\mathcal{L}_{\lambda_1, \dots, \lambda_n}$ and $T_{loc, \lambda_1, \dots, \lambda_n}^{m,*}$.

We begin studying the operator $T_{loc, \lambda_1, \dots, \lambda_n}^{m,*}$. For every $j = (j_1, \dots, j_n) \in \mathbb{Z}^n$, the dyadic cube Q_j is defined by

$$Q_j = \{y = (y_1, \dots, y_n) \in (0, \infty)^n : 2^{j_i} \leq y_i < 2^{j_i+1}, i = 1, \dots, n\},$$

and the cube $\tilde{Q}_j$ is given by

$$\tilde{Q}_j = \{y = (y_1, \dots, y_n) \in (0, \infty)^n : 2^{j_i-1} \leq y_i < 2^{j_i+2}, i = 1, \dots, n\}.$$

It is clear that if $j \in \mathbb{Z}^n$, $x \in Q_j$ and $y \in L(x)$, then $y \in \tilde{Q}_j$. We can write

$$\begin{aligned}
&\int_{L(x), |x-y| > \varepsilon} f(y) \prod_{l=1}^n (x_l y_l)^{-\lambda_l} H^\phi(x, y) \prod_{l=1}^n y_l^{2\lambda_l} dy = \int_{\tilde{Q}_j, |x-y| > \varepsilon} f(y) \prod_{l=1}^n \left(\frac{y_l}{x_l} \right)^{\lambda_l} H^\phi(x, y) dy \\
(41) \quad &- \int_{\tilde{Q}_j \setminus L(x), |x-y| > \varepsilon} f(y) \prod_{l=1}^n \left(\frac{y_l}{x_l} \right)^{\lambda_l} H^\phi(x, y) dy, \quad x \in Q_j, j \in \mathbb{Z}^n \text{ and } \varepsilon > 0.
\end{aligned}$$

Let $j = (j_1, \dots, j_n) \in \mathbb{Z}^n$. It has $\tilde{Q}_j \setminus L(x) = \bigcup_{i=1}^n (\tilde{Q}_{j,i}^+ \cup \tilde{Q}_{j,i}^-)$ where

$$\tilde{Q}_{j,i}^+ = \{y = (y_1, \dots, y_n) \in (0, \infty)^n : 2^{j_i-1} \leq y_l < 2^{j_i+2}, l = 1, \dots, n; l \neq i; 2x_i < y_i < 2^{j_i+2}\}$$

and

$$\tilde{Q}_{j,i}^- = \{y = (y_1, \dots, y_n) \in (0, \infty)^n : 2^{j_i-1} \leq y_l < 2^{j_i+2}, l = 1, \dots, n; l \neq i; 2^{j_i-1} < y_i < x_i/2\}$$

for $i = 1, \dots, n$.

We have that

$$\begin{aligned}
|H^\phi(x, y)| &\leq \int_0^\infty |\phi(t)| \left| \frac{\partial}{\partial t} \left(\frac{e^{-|x-y|^2/4t}}{t^{n/2}} \right) \right| dt \leq C \int_0^\infty \frac{e^{-|x-y|^2/8t}}{t^{n/2+1}} dt \\
(42) \quad &\leq C \frac{1}{|x-y|^n} \int_0^\infty \frac{e^{-1/u}}{u^{n/2+1}} du \leq C \frac{1}{|x-y|^n}, \quad x, y \in (0, \infty)^n.
\end{aligned}$$

By (42), for every $\varepsilon > 0$, we get

$$\begin{aligned}
& \left| \int_{\tilde{Q}_j \setminus L(x), |x-y| > \varepsilon} f(y) \prod_{l=1}^n \left(\frac{y_l}{x_l} \right)^{\lambda_l} H^\phi(x, y) dy \right| \leq \int_{\tilde{Q}_j \setminus L(x)} |f(y)| \prod_{l=1}^n \left(\frac{y_l}{x_l} \right)^{\lambda_l} |H^\phi(x, y)| dy \\
& \leq \sum_{i=1}^n \left(\int_{\tilde{Q}_{j,i}^+} |f(y)| \prod_{l=1}^n \left(\frac{y_l}{x_l} \right)^{\lambda_l} |H^\phi(x, y)| dy + \int_{\tilde{Q}_{j,i}^-} |f(y)| \prod_{l=1}^n \left(\frac{y_l}{x_l} \right)^{\lambda_l} |H^\phi(x, y)| dy \right) \\
& \leq C \sum_{i=1}^n \int_{\tilde{Q}_{j,i}^+ \cup \tilde{Q}_{j,i}^-} \frac{|f(y)|}{(x_i^2 + \sum_{l=1, l \neq i}^n (x_l - y_l)^2)^{n/2}} dy \\
& \leq C \sum_{i=1}^n \int_{\tilde{Q}_{j,i}^+ \cup \tilde{Q}_{j,i}^-} \frac{|f(y)|}{(2^{2j_i} + \sum_{l=1, l \neq i}^n (x_l - y_l)^2)^{n/2}} dy, \quad x \in Q_j.
\end{aligned}$$

Then, for each $x \in Q_j$,

$$(43) \quad \sup_{\varepsilon > 0} \left| \int_{\substack{\tilde{Q}_j \setminus L(x) \\ |x-y| > \varepsilon}} f(y) \prod_{l=1}^n \left(\frac{y_l}{x_l} \right)^{\lambda_l} H^\phi(x, y) dy \right| \leq C \sum_{i=1}^n \int_{\tilde{Q}_{j,i}^+ \cup \tilde{Q}_{j,i}^-} \frac{|f(y)|}{(2^{2j_i} + \sum_{\substack{l=1 \\ l \neq i}}^n (x_l - y_l)^2)^{n/2}} dy.$$

For every $i = 1, \dots, n$, we define

$$f_{j,i}(\bar{y}_i) = \int_{2^{j_i-1}}^{2^{j_i+2}} |f(y)| dy \chi_{\prod_{l=1, l \neq i}^n [2^{j_l-1}, 2^{j_l+2}]}(\bar{y}_i), \quad \bar{y}_i = (y_1, \dots, y_{i-1}, y_{i+1}, \dots, y_n) \in (0, \infty)^{n-1}.$$

From (43) it follows that

$$\begin{aligned}
& \sup_{\varepsilon > 0} \left| \int_{\substack{\tilde{Q}_j \setminus L(x) \\ |x-y| > \varepsilon}} f(y) \prod_{l=1}^n \left(\frac{y_l}{x_l} \right)^{\lambda_l} H^\phi(x, y) dy \right| \leq C \sum_{i=1}^n \int_{(0, \infty)^{n-1}} \frac{f_{j,i}(\bar{y}_i)}{(2^{2j_i} + |\bar{x}_i - \bar{y}_i|^2)^{n/2}} d\bar{y}_i \\
& \leq C \sum_{i=1}^n \left(\sum_{k=0}^{\infty} \int_{\substack{(0, \infty)^{n-1} \\ 2^{k+j_i} < |\bar{y}_i - \bar{x}_i| < 2^{k+j_i+1}}} \frac{f_{j,i}(\bar{y}_i)}{(2^{2j_i} + |\bar{x}_i - \bar{y}_i|^2)^{n/2}} d\bar{y}_i \right. \\
& \quad \left. + \int_{\substack{(0, \infty)^{n-1} \\ |\bar{y}_i - \bar{x}_i| < 2^{j_i}}} \frac{f_{j,i}(\bar{y}_i)}{(2^{2j_i} + |\bar{x}_i - \bar{y}_i|^2)^{n/2}} d\bar{y}_i \right) \\
& \leq C \sum_{i=1}^n \left(\sum_{k=0}^{\infty} \frac{1}{2^{nj_i}(1+2^{2k})^{n/2}} \int_{|\bar{y}_i - \bar{x}_i| < 2^{k+j_i+1}} f_{j,i}(\bar{y}_i) d\bar{y}_i + \frac{1}{2^{nj_i}} \int_{|\bar{y}_i - \bar{x}_i| < 2^{j_i}} f_{j,i}(\bar{y}_i) d\bar{y}_i \right) \\
& \leq C \sum_{i=1}^n \frac{1}{2^{j_i}} M_{n-1}(f_{j,i})(\bar{x}_i), \quad x \in Q_j.
\end{aligned}$$

Here, M_{n-1} represents the Hardy-Littlewood maximal function on $\mathbb{R}^{n-1}$. We recall that M_{n-1} is a bounded operator from $L^p(\mathbb{R}^{n-1})$ into itself, for every $1 < p < \infty$, and from $L^1(\mathbb{R}^{n-1})$ into $L^{1,\infty}(\mathbb{R}^{n-1})$.

On the other hand, it is known that the maximal operator $T^{m,*}$ defined by

$$T^{m,*}(f)(x) = \sup_{\varepsilon > 0} \left| \int_{|x-y| > \varepsilon} f(y) H^\phi(x, y) dy \right|, \quad x \in \mathbb{R}^n,$$

is bounded from $L^p(\mathbb{R}^n, dx)$ into itself, for every $1 < p < \infty$, and from $L^1(\mathbb{R}^n, dx)$ into $L^{1,\infty}(\mathbb{R}^n, dx)$.

We denotes by $m_{\lambda_1, \dots, \lambda_n}^{(n)}$ the positive measure given by $m_{\lambda_1, \dots, \lambda_n}^{(n)}(E) = \int_E \prod_{j=1}^n x_j^{2\lambda_j} dx$, for every Lebesgue measurable set E .

Let $\gamma > 0$. We have that,

$$\begin{aligned}
& m_{\lambda_1, \dots, \lambda_n}^{(n)}(\{x \in (0, \infty)^n : T_{loc, \lambda_1, \dots, \lambda_n}^{m, *} (f)(x) > \gamma\}) \\
&= \sum_{j \in \mathbb{Z}^n} m_{\lambda_1, \dots, \lambda_n}^{(n)}(\{x \in Q_j : T_{loc, \lambda_1, \dots, \lambda_n}^{m, *} (f)(x) > \gamma\}) \\
&\leq \sum_{j \in \mathbb{Z}^n} \left(m_{\lambda_1, \dots, \lambda_n}^{(n)}(\{x \in Q_j : T^{m, *}(\prod_{l=1}^n (y_l/x_l)^{\lambda_l} f \chi_{\tilde{Q}_j})(x) > C\gamma\}) \right. \\
&\quad \left. + m_{\lambda_1, \dots, \lambda_n}^{(n)}\left(\left\{x \in Q_j : \sup_{\varepsilon > 0} \left| \int_{\tilde{Q}_j \setminus L(x)} f(y) \prod_{l=1}^n \left(\frac{y_l}{x_l}\right)^{\lambda_l} H^\phi(x, y) dy \right| > C\gamma \right\}\right) \right) \\
&\leq \sum_{j \in \mathbb{Z}^n} \left(\frac{C}{\gamma} \prod_{i=1}^n 2^{2j_i \lambda_i} \|f \chi_{\tilde{Q}_j}\|_{L^1(\mathbb{R}^n)} + \sum_{i=1}^n m_{\lambda_1, \dots, \lambda_n}^{(n)}(\{x \in \tilde{Q}_j : M_{n-1}(f_{j,i})(\bar{x}_i) > 2^{j_i} C\gamma\}) \right) \\
&\leq \sum_{j \in \mathbb{Z}^n} \left(\frac{C}{\gamma} \|f \chi_{\tilde{Q}_j}\|_{L^1((0, \infty)^n, \prod_{i=1}^n x_i^{2\lambda_i} dx)} \right. \\
&\quad \left. + \sum_{i=1}^n 2^{(2\lambda_i+1)j_i} m_{\lambda_1, \dots, \lambda_{i-1}, \lambda_{i+1}, \dots, \lambda_n}^{(n-1)}(\{\bar{x}_i \in \prod_{l=1, l \neq i}^n [2^{j_l}, 2^{j_l+1}) : M_{n-1}(f_{j,i})(\bar{x}_i) > 2^{j_i} C\gamma\}) \right) \\
&\leq \frac{C}{\gamma} \sum_{j \in \mathbb{Z}^n} \left(\|f \chi_{\tilde{Q}_j}\|_{L^1((0, \infty)^n, \prod_{i=1}^n x_i^{2\lambda_i} dx)} + \sum_{i=1}^n \|f_{j,i}\|_{L^1(\mathbb{R}^{n-1})} \prod_{l=1}^n 2^{2j_l \lambda_l} \right) \\
&\leq \frac{C}{\gamma} \sum_{j \in \mathbb{Z}^n} \|f \chi_{\tilde{Q}_j}\|_{L^1((0, \infty)^n, \prod_{i=1}^n x_i^{2\lambda_i} dx)} \leq \frac{C}{\gamma} \|f\|_{L^1((0, \infty)^n, \prod_{i=1}^n x_i^{2\lambda_i} dx)}, \quad f \in L^1((0, \infty)^n, \prod_{i=1}^n x_i^{2\lambda_i} dx).
\end{aligned}$$

Hence, $T_{loc, \lambda_1, \dots, \lambda_n}^{m, *}$ is bounded from $L^1((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$ into $L^{1, \infty}((0, \infty)^n, \prod_{i=1}^n x_i^{2\lambda_i} dx)$. Also, if $1 < p < \infty$, we have

$$\begin{aligned}
& \|T_{loc, \lambda_1, \dots, \lambda_n}^{m, *} (f)\|_{L^p((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)}^p \leq \sum_{j \in \mathbb{Z}^n} \|T_{loc, \lambda_1, \dots, \lambda_n}^{m, *} (f) \chi_{Q_j}\|_{L^p(\mathbb{R}^n, dx)}^p \prod_{l=1}^n 2^{2\lambda_l j_l} \\
&\leq C \sum_{j \in \mathbb{Z}^n} \left(\int_{Q_j} |T^{m, *}(\prod_{l=1}^n (y_l/x_l)^{\lambda_l} f \chi_{\tilde{Q}_j})(x)|^p dx + \sum_{i=1}^n \frac{1}{2^{j_i p}} \int_{Q_j} |M_{n-1}(f_{j,i})(\bar{x}_i)|^p d\bar{x}_i dx_i \right) \prod_{l=1}^n 2^{2\lambda_l j_l} \\
&\leq C \sum_{j \in \mathbb{Z}^n} \left(\int_{\tilde{Q}_j} |f(x)|^p dx + \sum_{i=1}^n \frac{1}{2^{j_i p}} \int_{2^{j_i}}^{2^{j_i+1}} \|f_{j,i}\|_{L^p(\mathbb{R}^{n-1})}^p dx_i \right) \prod_{l=1}^n 2^{2\lambda_l j_l} \\
&\leq C \sum_{j \in \mathbb{Z}^n} \int_{\tilde{Q}_j} |f(x)|^p dx \prod_{l=1}^n 2^{2\lambda_l j_l} \leq C \|f\|_{L^p((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)}^p, \quad f \in L^p((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx).
\end{aligned}$$

It is proved that $T_{loc, \lambda_1, \dots, \lambda_n}^{m, *}$ is bounded from $L^p((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$ into itself, for every $1 < p < \infty$.

We now analyze the operator $\mathcal{L}_{\lambda_1, \dots, \lambda_n}$. Let $\alpha > -1/2$. From (5) and (37) we deduce that, if $t, u, v \in (0, \infty)$ and $uv/t > 1$,

$$\begin{aligned}
\frac{\partial}{\partial t} W_t^\alpha(u, v) &= \left\{ -\frac{\alpha + 1/2}{t} \left(\frac{uv}{2t} \right)^{1/2} I_{\alpha-1/2} \left(\frac{uv}{2t} \right) - \frac{uv}{2t^2} \left(\frac{uv}{2t} \right)^{1/2} I_{\alpha+1/2} \left(\frac{uv}{2t} \right) \right. \\
&\quad \left. + \frac{u^2 + v^2}{4t^2} \left(\frac{uv}{2t} \right)^{1/2} I_{\alpha-1/2} \left(\frac{uv}{2t} \right) \right\} \frac{(uv)^{-\alpha}}{\sqrt{2t}} e^{-(u^2+v^2)/4t} \\
&= \left\{ -\frac{\alpha + 1/2}{t} \left(1 + \mathcal{O} \left(\frac{t}{uv} \right) \right) - \frac{uv}{2t^2} \left(1 - \frac{4(\alpha + 1/2)^2 - 1}{4} \frac{t}{uv} + \mathcal{O} \left(\left(\frac{t}{uv} \right)^2 \right) \right) \right. \\
&\quad \left. + \frac{u^2 + v^2}{4t^2} \left(1 - \frac{4(\alpha + 1/2)^2 - 1}{4} \frac{t}{uv} + \mathcal{O} \left(\left(\frac{t}{uv} \right)^2 \right) \right) \right\} \frac{(uv)^{-\alpha}}{\sqrt{4\pi t}} e^{-(u-v)^2/4t} \\
&= \left\{ \frac{(u-v)^2}{4t^2} - \frac{1}{2t} + \frac{(u-v)^2}{4tuv} + \mathcal{O} \left(\frac{1}{uv} \right) + \mathcal{O} \left(\frac{(u-v)^2}{(uv)^2} \right) \right\} \frac{(uv)^{-\alpha}}{\sqrt{4\pi t}} e^{-(u-v)^2/4t} \\
(44) \quad &= \frac{\partial}{\partial t} \left(\frac{e^{(u-v)^2/4t}}{\sqrt{4\pi t}} \right) (uv)^{-\alpha} + \mathcal{O} \left(\frac{(uv)^{-\alpha}}{t^{1/2}} e^{-(u-v)^2/8t} \left(\frac{1}{uv} + \frac{(u-v)^2}{(uv)^2} \right) \right).
\end{aligned}$$

By proceeding in a similar way we can see that

$$(45) \quad \left| W_t^\alpha(u, v) - (uv)^{-\alpha} \frac{e^{-(u-v)^2/4t}}{\sqrt{4\pi t}} \right| \leq C(uv)^{-\alpha-1} \sqrt{t} e^{-(u-v)^2/4t}, \quad t, u, v \in (0, \infty) \text{ and } \frac{uv}{t} > 1.$$

We have that

$$\begin{aligned}
|\mathcal{L}_{\lambda_1, \dots, \lambda_n}(f)(x)| &\leq \int_{L(x)} |f(y)| |K_{\lambda_1, \dots, \lambda_n}^\phi(x, y) - \prod_{j=1}^n (x_j y_j)^{-\lambda_j} H^\phi(x, y)| \prod_{j=1}^n y_j^{2\lambda_j} dy \\
&\leq \int_{L(x)} |f(y)| \int_0^\infty |\phi(t)| \left| \frac{\partial}{\partial t} \left(\prod_{j=1}^n W_t^{\lambda_j}(x_j, y_j) \right) - \prod_{j=1}^n (x_j y_j)^{-\lambda_j} \frac{\partial}{\partial t} \left(\prod_{j=1}^n \frac{e^{-(x_j-y_j)^2/4t}}{\sqrt{4\pi t}} \right) \right| dt \prod_{j=1}^n y_j^{2\lambda_j} dy \\
&\leq C \sum_{i=1}^n \int_{L(x)} |f(y)| \int_0^\infty \left| \prod_{\substack{j=1 \\ j \neq i}}^n W_t^{\lambda_j}(x_j, y_j) \frac{\partial}{\partial t} W_t^{\lambda_i}(x_i, y_i) \right. \\
&\quad \left. - \prod_{j=1}^n (x_j y_j)^{-\lambda_j} \prod_{\substack{j=1 \\ j \neq i}}^n \frac{e^{-(x_j-y_j)^2/4t}}{\sqrt{4\pi t}} \frac{\partial}{\partial t} \left(\frac{e^{-(x_i-y_i)^2/4t}}{\sqrt{4\pi t}} \right) \right| dt \prod_{j=1}^n y_j^{2\lambda_j} dy, \quad x \in (0, \infty)^n.
\end{aligned}$$

We analyze the operator defined by the first summand. The other ones can be studied similarly. It follows that, for each $x \in (0, \infty)^n$,

$$\begin{aligned}
\mathcal{L}_{\lambda_1, \dots, \lambda_n}^1(f)(x) &= \int_{L(x)} |f(y)| \int_0^\infty \left| \prod_{j=2}^n W_t^{\lambda_j}(x_j, y_j) \frac{\partial}{\partial t} W_t^{\lambda_1}(x_1, y_1) \right. \\
&\quad \left. - \prod_{j=1}^n (x_j y_j)^{-\lambda_j} \prod_{j=2}^n \frac{e^{-(x_j-y_j)^2/4t}}{\sqrt{4\pi t}} \frac{\partial}{\partial t} \left(\frac{e^{-(x_1-y_1)^2/4t}}{\sqrt{4\pi t}} \right) \right| dt \prod_{j=1}^n y_j^{2\lambda_j} dy \\
&\leq \int_{L(x)} |f(y)| \int_0^\infty \left| \frac{\partial}{\partial t} W_t^{\lambda_1}(x_1, y_1) - (x_1 y_1)^{-\lambda_1} \frac{\partial}{\partial t} \left(\frac{e^{-(x_1-y_1)^2/4t}}{\sqrt{4\pi t}} \right) \right| \prod_{j=2}^n W_t^{\lambda_j}(x_j, y_j) dt \\
&\quad \times \prod_{j=1}^n y_j^{2\lambda_j} dy + \sum_{i=2}^n \int_{L(x)} |f(y)| \int_0^\infty (x_1 y_1)^{-\lambda_1} \left| \frac{\partial}{\partial t} \left(\frac{e^{-(x_1-y_1)^2/4t}}{\sqrt{4\pi t}} \right) \right| \prod_{j=2}^{i-1} \left(\frac{e^{-(x_j-y_j)^2/4t}}{\sqrt{4\pi t}} \right) \\
(46) \quad &\quad \times (x_j y_j)^{-\lambda_j} \left| W_t^{\lambda_i}(x_i, y_i) - (x_i y_i)^{-\lambda_i} \frac{e^{-(x_i-y_i)^2/4t}}{\sqrt{4\pi t}} \right| \prod_{j=i+1}^n W_t^{\lambda_j}(x_j, y_j) dt \prod_{j=1}^n y_j^{2\lambda_j} dy.
\end{aligned}$$

We now use (4), (5), (37), and (44) to obtain the following

$$\begin{aligned}
\mathcal{L}_{\lambda_1, \dots, \lambda_n}^{1,1}(f)(x) &= \int_{L(x)} |f(y)| \int_0^\infty \left| \frac{\partial}{\partial t} W_t^{\lambda_1}(x_1, y_1) - (x_1 y_1)^{-\lambda_1} \frac{\partial}{\partial t} \left(\frac{e^{-\frac{(x_1 - y_1)^2}{4t}}}{\sqrt{4\pi t}} \right) \right| \\
&\quad \times \prod_{j=2}^n W_t^{\lambda_j}(x_j, y_j) dt \prod_{j=1}^n y_j^{2\lambda_j} dy \\
&\leq \int_{L(x)} |f(y)| \left\{ \int_0^{x_1 y_1} \frac{(x_1 y_1)^{-\lambda_1 - 1}}{t^{1/2}} e^{-\frac{(x_1 - y_1)^2}{10t}} \prod_{j=2}^n \left(\frac{1}{x_j^{2\lambda_j + 1}} + (x_j y_j)^{-\lambda_j} \frac{e^{-\frac{(x_j - y_j)^2}{4t}}}{\sqrt{t}} \right) dt \right. \\
(47) \quad &\quad \left. + \int_{x_1 y_1}^\infty \left(\frac{1}{t^{\lambda_1 + \frac{3}{2}}} + (x_1 y_1)^{-\lambda_1} \frac{1}{t^{\frac{3}{2}}} \right) \prod_{j=2}^n \left(\frac{1}{x_j^{2\lambda_j + 1}} + (x_j y_j)^{-\lambda_j} \frac{e^{-\frac{(x_j - y_j)^2}{4t}}}{\sqrt{t}} \right) dt \right\} \prod_{j=1}^n y_j^{2\lambda_j} dy
\end{aligned}$$

for every $x \in (0, \infty)^n$.

Then, the operator $\mathcal{L}_{\lambda_1, \dots, \lambda_n}^{1,1}$ is controlled by operators of the following type:

$$\Lambda_{\lambda_1, \dots, \lambda_n}^l(g)(x) = \sup_{t>0} \left| \int_{\frac{x_{l+1}}{2}}^{2x_{l+1}} \dots \int_{\frac{x_n}{2}}^{2x_n} \prod_{j=l+1}^n x_j^{-2\lambda_j - 1} \int_{\frac{x_1}{2}}^{2x_1} \dots \int_{\frac{x_l}{2}}^{2x_l} \prod_{j=1}^l (x_j y_j)^{-\lambda_j} \frac{e^{-\frac{(x_j - y_j)^2}{10t}}}{\sqrt{t}} g(y) \prod_{j=1}^n y_j^{2\lambda_j} dy \right|,$$

where $0 \leq l \leq n$, $l \in \mathbb{N}$. For each $0 \leq l \leq n$, $l \in \mathbb{N}$, $\Lambda_{\lambda_1, \dots, \lambda_n}^l$ is bounded operator from $L^p((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$ into itself, for every $1 < p < \infty$, and from $L^1((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$ into $L^{1,\infty}((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$. Indeed, let $k \in \mathbb{N}$. We consider the maximal operators

$$\Omega_{\beta_1, \dots, \beta_k}(g)(x) = \sup_{t>0} \left| \int_{\frac{x_1}{2}}^{2x_1} \dots \int_{\frac{x_k}{2}}^{2x_k} \prod_{j=1}^k (x_j y_j)^{-\beta_j} \frac{e^{-\frac{(x_j - y_j)^2}{4t}}}{\sqrt{t}} g(y) \prod_{j=1}^k y_j^{2\beta_j} dy \right|, \quad x \in (0, \infty)^k,$$

where $\beta_j \in \mathbb{R}$, $j = 1, \dots, k$, and

$$M_k(g)(x) = \sup_{t>0} \left| \int_{(0, \infty)^k} \frac{e^{-|x-y|^2/4t}}{t^{k/2}} g(y) dy \right|, \quad x \in (0, \infty)^k.$$

For every $j = (j_1, \dots, j_k) \in \mathbb{Z}^k$ we define

$$Q_j = \{y = (y_1, \dots, y_k) \in (0, \infty)^k : 2^{j_i} \leq y_i < 2^{j_i+1}, i = 1, \dots, k\},$$

and

$$\tilde{Q}_j = \{y = (y_1, \dots, y_k) \in (0, \infty)^k : 2^{j_i-1} \leq y_i < 2^{j_i+2}, i = 1, \dots, k\}.$$

Assume that $\gamma > 0$. Since, as it well known, the operator M_k is bounded from $L^1((0, \infty)^k, dx)$ into $L^{1,\infty}((0, \infty)^k, dx)$, we have

$$\begin{aligned}
m_{\beta_1, \dots, \beta_k}^{(k)}(\{x \in (0, \infty)^k : \Omega_{\beta_1, \dots, \beta_k}(g)(x) > \lambda\}) &= \sum_{j \in \mathbb{Z}^k} m_{\beta_1, \dots, \beta_k}^{(k)}(\{x \in Q_j : \Omega_{\beta_1, \dots, \beta_k}(g)(x) > \lambda\}) \\
&\leq \sum_{j \in \mathbb{Z}^k} 2^{2 \sum_{i=1}^k \beta_i j_i} m_{0, \dots, 0}^{(k)}(\{x \in \mathbb{R}^k : M_k(|g| \chi_{\tilde{Q}_j})(x) > C\gamma\}) \\
&\leq \frac{C}{\gamma} \sum_{j \in \mathbb{Z}^k} 2^{2 \sum_{i=1}^k \beta_i j_i} \int_{\tilde{Q}_j} |g(y)| dy \\
&\leq \frac{C}{\gamma} \int_{(0, \infty)^k} |g(y)| \prod_{j=1}^k y_j^{2\beta_j} dy, \quad g \in L^1((0, \infty)^k, \prod_{j=1}^k x_j^{2\beta_j} dx).
\end{aligned}$$

Also, M_k is bounded from $L^p((0, \infty)^k, dx)$ into itself, for every $1 < p < \infty$, and it has

$$\begin{aligned}
\int_{(0, \infty)^k} |\Omega_{\beta_1, \dots, \beta_k}(g)(x)|^p \prod_{j=1}^k x_j^{2\beta_j} dx &= 2^{2 \sum_{i=1}^k \beta_i j_i} \sum_{j \in \mathbb{Z}^k} \int_{Q_j} |\Omega_{\beta_1, \dots, \beta_k}(g)(x)|^p dx \\
&\leq C \sum_{j \in \mathbb{Z}^k} 2^{2 \sum_{i=1}^k \beta_i j_i} \int_{\mathbb{R}^k} |M_k(|g| \chi_{\tilde{Q}_j})(x)|^p dx \\
&\leq C 2^{2 \sum_{i=1}^k \beta_i j_i} \sum_{j \in \mathbb{Z}^k} \int_{\tilde{Q}_j} |g(y)|^p dy \\
&\leq C \int_{(0, \infty)^k} |g(y)|^p \prod_{j=1}^k y_j^{2\beta_j} dy, \quad g \in L^p((0, \infty)^k, \prod_{j=1}^k x_j^{2\beta_j} dx),
\end{aligned}$$

and $\Omega_{\beta_1, \dots, \beta_k}$ is bounded from $L^p((0, \infty)^k, \prod_{j=1}^k x_j^{2\beta_j} dx)$ into itself, for every $1 < p < \infty$.

On the other hand, it is not hard to see that, for every $k \in \mathbb{N}$ and $\beta_j \in \mathbb{R}$, $j = 1, \dots, k$, the operator $Z_{\beta_1, \dots, \beta_k}$ defined by

$$Z_{\beta_1, \dots, \beta_k}(g)(x) = \prod_{j=1}^k x_j^{-2\beta_j-1} \int_{x_1/2}^{2x_1} \dots \int_{x_k/2}^{2x_k} g(y) \prod_{j=1}^k z_j^{2\beta_j} dy, \quad x \in (0, \infty)^k,$$

is bounded from $L^p((0, \infty)^k, \prod_{j=1}^k x_j^{2\beta_j} dx)$ into itself, for every $1 \leq p < \infty$.

Then, the L^p -boundedness properties of operators $\Omega_{\beta_1, \dots, \beta_k}$ and $Z_{\beta_1, \dots, \beta_k}$, $\beta_j > -1/2$, $j = 1, \dots, k$, $k \in \mathbb{N}$, allow us, by using [4, Proposition 1], to conclude that, for every $m \in \mathbb{N}$, the operator $\Lambda_{\lambda_1, \dots, \lambda_n}^m$ is bounded from $L^p((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$ into itself, for every $1 < p < \infty$, and from $L^1((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$ into $L^{1, \infty}((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$. Hence, the operator $\mathcal{L}_{\lambda_1, \dots, \lambda_n}^{1,1}$ is bounded from $L^p((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$ into itself, for every $1 < p < \infty$, and from $L^1((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$ into $L^{1, \infty}((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$.

For every $i = 2, \dots, n$, it has

$$\begin{aligned}
\mathcal{L}_{\lambda_1, \dots, \lambda_n}^{1,i}(f)(x) &= \int_{L(x)} |f(y)| \int_0^\infty (x_1 y_1)^{-\lambda_1} \left| \frac{\partial}{\partial t} \left(\frac{e^{-\frac{(x_1 - y_1)^2}{4t}}}{\sqrt{4\pi t}} \right) \right| \prod_{j=2}^{i-1} \left(\frac{e^{-\frac{(x_j - y_j)^2}{4t}}}{\sqrt{4\pi t}} \right) \\
&\quad \times (x_j y_j)^{-\lambda_j} \left| W_t^{\lambda_i}(x_i, y_i) - (x_i y_i)^{-\lambda_i} \frac{e^{-\frac{(x_i - y_i)^2}{4t}}}{\sqrt{4\pi t}} \right| \prod_{j=i+1}^n W_t^{\lambda_j}(x_j, y_j) dt \prod_{j=1}^n y_j^{2\lambda_j} dy \\
(48) \quad &\leq C \int_{L(x)} |f(y)| \left\{ \int_0^{x_i y_i} \frac{e^{-\sum_{j=1}^{i-1} \frac{(x_j - y_j)^2}{4t}}}{t^{\frac{i+1}{2}}} \prod_{j=1}^{i-1} (x_j y_j)^{-\lambda_j} (x_i y_i)^{-\lambda_i - 1} \sqrt{t} e^{-\frac{(x_i - y_i)^2}{4t}} \right. \\
&\quad \times \prod_{j=i+1}^n \left(\frac{1}{x_j^{2\lambda_j+1}} + (x_j y_j)^{-\lambda_j} \frac{e^{-\frac{(x_j - y_j)^2}{4t}}}{\sqrt{t}} \right) dt + \int_{x_i y_i}^\infty \prod_{j=1}^{i-1} (x_j y_j)^{-\lambda_j} \frac{e^{-\sum_{j=1}^{i-1} \frac{(x_j - y_j)^2}{4t}}}{t^{\frac{i+1}{2}}} \left(\frac{1}{t^{\lambda_i + \frac{1}{2}}} \right. \\
&\quad \left. \left. + (x_i y_i)^{-\lambda_i} \frac{e^{-\frac{(x_i - y_i)^2}{4t}}}{\sqrt{t}} \right) \prod_{j=i+1}^n \left(\frac{1}{x_j^{2\lambda_j+1}} + (x_j y_j)^{-\lambda_j} \frac{e^{-\frac{(x_j - y_j)^2}{4t}}}{\sqrt{t}} \right) dt \right\} \prod_{j=1}^n y_j^{2\lambda_j} dy, \quad x \in (0, \infty)^n.
\end{aligned}$$

Then, the operator $\mathcal{L}_{\lambda_1, \dots, \lambda_n}^{1,i}$ can be controlled by operator of type $\Lambda_{\lambda_1, \dots, \lambda_n}^m$, $m \in \mathbb{N}$. Thus, we conclude that $\mathcal{L}_{\lambda_1, \dots, \lambda_n}^{1,i}$ is bounded from $L^p((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$ into itself, for every $1 < p < \infty$, and from $L^1((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$ into $L^{1,\infty}((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$.

From the assertions proved in (47) and (48), $i = 2, \dots, n$, we deduce that the operator $\mathcal{L}_{\lambda_1, \dots, \lambda_n}^1$ is bounded from $L^p((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$ into itself, for every $1 < p < \infty$, and from $L^1((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$ into $L^{1,\infty}((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$.

Thus we conclude that the operator $\mathcal{L}_{\lambda_1, \dots, \lambda_n}$ is bounded from $L^p((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$ into itself, for every $1 < p < \infty$, and from $L^1((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$ into $L^{1,\infty}((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$.

Finally we study the operator $\mathcal{G}_{\lambda_1, \dots, \lambda_n}^\phi$. It is clear that $\mathcal{G}_{\lambda_1, \dots, \lambda_n}^\phi$ can be decomposed in a sum of operators like the following one

$$\begin{aligned}
 S_{\lambda_1, \dots, \lambda_n}^{k,l}(g)(x) &= \int_0^{\frac{x_1}{2}} \cdots \int_0^{\frac{x_k}{2}} \int_{\frac{x_{k+1}}{2}}^{2x_{k+1}} \cdots \int_{\frac{x_l}{2}}^{2x_l} \int_{2x_{l+1}}^\infty \cdots \int_{2x_n}^\infty g(y) K_{\lambda_1, \dots, \lambda_n}^\phi(x, y) \prod_{j=1}^n y_j^{2\lambda_j} dy \\
 &= \sum_{i=1}^n \int_0^{\frac{x_1}{2}} \cdots \int_0^{\frac{x_k}{2}} \int_{\frac{x_{k+1}}{2}}^{2x_{k+1}} \cdots \int_{\frac{x_l}{2}}^{2x_l} \int_{2x_{l+1}}^\infty \cdots \int_{2x_n}^\infty g(y) K_{\lambda_1, \dots, \lambda_n}^{\phi,i}(x, y) \prod_{j=1}^n y_j^{2\lambda_j} dy \\
 (49) \quad &= \sum_{i=1}^n S_{\lambda_1, \dots, \lambda_n}^{k,l,i}(g)(x), \quad x \in (0, \infty)^n,
 \end{aligned}$$

where $0 \leq k \leq l \leq n$, $k \neq 0$ or $l \neq n$, and

$$K_{\lambda_1, \dots, \lambda_n}^{\phi,i}(x, y) = \int_0^\infty \prod_{j=1}^{i-1} W_t^{\lambda_j}(x_j, y_j) \frac{\partial}{\partial t} W_t^{\lambda_i}(x_i, y_i) \prod_{j=i+1}^n W_t^{\lambda_j}(x_j, y_j) \phi(t) dt, \quad x, y \in (0, \infty)^n,$$

for every $i = 1, \dots, n$. From (5) we deduce that, if $\alpha > -1/2$,

$$\begin{aligned}
 |W_t^\alpha(u, v)| &\leq C \frac{(uv)^{-\alpha} e^{-v^2/16t}}{\sqrt{t}} \leq C (uv)^{-\alpha-1/2} \left(\frac{uv}{t}\right)^{1/2} e^{-v^2/16t} \\
 (50) \quad &\leq C t^{-\alpha-1/2} \frac{v}{\sqrt{t}} e^{-v^2/16t} \leq \frac{C}{t^{\alpha+1/2}} e^{-v^2/20t}, \quad t, u, v \in (0, \infty), \frac{uv}{t} > 1 \text{ and } 2u < v < \infty.
 \end{aligned}$$

By (4) and (50) we conclude that for $\alpha > -1/2$

$$(51) \quad |W_t^\alpha(u, v)| \leq \frac{C}{t^{\alpha+1/2}} e^{-v^2/20t}, \quad t, u, v \in (0, \infty), \text{ and } 2u < v < \infty.$$

From (44) it follows, if $\alpha > -1/2$,

$$\begin{aligned}
 \left| \frac{\partial}{\partial t} W_t^\alpha(u, v) \right| &\leq C e^{-v^2/16t} \left(\frac{1}{t^{3/2}} (uv)^{-\alpha} + \frac{(uv)^{-\alpha-1}}{t^{1/2}} \right) \\
 (52) \quad &\leq C \frac{e^{-v^2/16t}}{t^{3/2}} (uv)^{-\alpha} \leq C \frac{e^{-v^2/20t}}{t^{\alpha+3/2}}, \quad t, u, v \in (0, \infty), \frac{uv}{t} > 1 \text{ and } 2u < v < \infty.
 \end{aligned}$$

By (37) and (52) we get for every $\alpha > -1/2$

$$(53) \quad \left| \frac{\partial}{\partial t} W_t^\alpha(u, v) \right| \leq C \frac{e^{-v^2/20t}}{t^{\alpha+3/2}}, \quad t, u, v \in (0, \infty), \quad 2u < v < \infty.$$

According to (51) and (53) we infer

$$\begin{aligned}
\left| S_{\lambda_1, \dots, \lambda_n}^{0,0,i}(g)(x) \right| &\leq C \int_{2x_1}^{\infty} \dots \int_{2x_n}^{\infty} |g(y)| \int_0^{\infty} \frac{e^{-\sum_{j=1}^n y_j^2/20t}}{t^{\sum_{j=1}^n (\lambda_j+1/2)+1}} dt \prod_{j=1}^n y_j^{2\lambda_j} dy \\
&\leq C \int_{2x_1}^{\infty} \dots \int_{2x_n}^{\infty} |g(y)| \frac{1}{\left(\sum_{j=1}^n y_j^2\right)^{\sum_{j=1}^n (\lambda_j+1/2)}} \prod_{j=1}^n y_j^{2\lambda_j} dy \\
&\leq C \int_{2x_1}^{\infty} \dots \int_{2x_n}^{\infty} |g(y)| \frac{1}{y_1 \dots y_n} dy, \quad x \in (0, \infty)^n \text{ and } i = 1, \dots, n.
\end{aligned}$$

It is not hard to see that, for every $k \in \mathbb{N}$ the operator

$$\mathcal{S}_k(g)(x) = \int_{2x_1}^{\infty} \dots \int_{2x_k}^{\infty} |g(y)| \frac{1}{y_1 \dots y_k} dy, \quad x \in (0, \infty)^k,$$

is bounded from $L^p((0, \infty)^k, \prod_{j=1}^k x_j^{2\beta_j} dx)$ into itself, for every $1 \leq p < \infty$ and $\beta_j > -1/2$, $j = 1, \dots, k$.

Hence the operator $S_{\lambda_1, \dots, \lambda_n}^{0,0,i}$ is bounded from $L^p((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$, for every $1 \leq p < \infty$ and $i = 1, \dots, n$.

Let $i, k \in \mathbb{N}$, $1 \leq i \leq k$, and $\beta_j > -1/2$, $j = 1, \dots, k$. We define the operator $\mathcal{H}_{\beta_1, \dots, \beta_k}$ by

$$\begin{aligned}
\mathcal{H}_{\beta_1, \dots, \beta_k}(g)(x) &= \int_0^{\frac{x_1}{2}} \dots \int_0^{\frac{x_k}{2}} |g(y)| \int_0^{\infty} \prod_{j=1}^{i-1} W_t^{\beta_j}(x_j, y_j) \left| \frac{\partial}{\partial t} W_t^{\beta_i}(x_i, y_i) \right| \\
&\quad \times \prod_{j=i+1}^k W_t^{\beta_j}(x_j, y_j) dt \prod_{j=1}^k y_j^{2\beta_j} dy, \quad x \in (0, \infty)^k.
\end{aligned}$$

By taking into account symmetries, (51), and (53), we get that

$$\begin{aligned}
|\mathcal{H}_{\beta_1, \dots, \beta_k}(g)(x)| &\leq C \int_0^{\frac{x_1}{2}} \dots \int_0^{\frac{x_k}{2}} |g(y)| \int_0^{\infty} \frac{e^{-\sum_{j=1}^k x_j^2/20t}}{t^{\sum_{j=1}^k (\beta_j+1/2)+1}} dt \prod_{j=1}^k y_j^{2\beta_j} dy \\
(54) \quad &\leq \frac{C}{\left(\sum_{j=1}^k x_j^2\right)^{\sum_{j=1}^k (\beta_j+1/2)}} \int_0^{\frac{x_1}{2}} \dots \int_0^{\frac{x_k}{2}} |g(y)| \prod_{j=1}^k y_j^{2\beta_j} dy.
\end{aligned}$$

The operator $H_{\beta_1, \dots, \beta_k}$ given by

$$H_{\beta_1, \dots, \beta_k}(g)(x) = \frac{1}{\left(\sum_{j=1}^k x_j^2\right)^{\sum_{j=1}^k (\beta_j+1/2)}} \int_0^{\frac{x_1}{2}} \dots \int_0^{\frac{x_k}{2}} g(y) \prod_{j=1}^k y_j^{2\beta_j} dy$$

is bounded from $L^p((0, \infty)^k, \prod_{j=1}^k x_j^{2\beta_j} dx)$ into itself, for every $1 < p < \infty$, and from $L^1((0, \infty)^k, \prod_{j=1}^k x_j^{2\beta_j} dx)$ into $L^{1,\infty}((0, \infty)^k, \prod_{j=1}^k x_j^{2\beta_j} dx)$. Indeed, assume that $\gamma > 0$. We have

$$\begin{aligned} & m_{\beta_1, \dots, \beta_k}^{(k)}(\{x \in (0, \infty)^k : |H_{\beta_1, \dots, \beta_k}(g)(x)| > \gamma\}) \\ & \leq m_{\beta_1, \dots, \beta_k}^{(k)}\left(\left\{x \in (0, \infty)^k : \sum_{j=1}^k x_j^2 \leq \left(\frac{1}{\gamma} \|g\|_{L^1((0, \infty)^k, \prod_{j=1}^k x_j^{2\beta_j} dx)}\right)^{\frac{1}{\sum_{j=1}^k (\beta_j + 1/2)}}\right\}\right) \\ & \leq m_{\beta_1, \dots, \beta_k}^{(k)}\left(\left\{x \in (0, \infty)^k : 0 \leq x_j \leq \left(\frac{1}{\gamma} \|g\|_{L^1((0, \infty)^k, \prod_{j=1}^k x_j^{2\beta_j} dx)}\right)^{\frac{1}{\sum_{j=1}^k (2\beta_j + 1)}}, j = 1, \dots, k\right\}\right) \\ & \leq \frac{C}{\gamma} \|g\|_{L^1((0, \infty)^k, \prod_{j=1}^k x_j^{2\beta_j} dx)}, \quad g \in L^1((0, \infty)^k, \prod_{j=1}^k x_j^{2\beta_j} dx). \end{aligned}$$

Hence, $H_{\beta_1, \dots, \beta_k}$ is bounded from $L^1((0, \infty)^k, \prod_{j=1}^k x_j^{2\beta_j} dx)$ into $L^{1,\infty}((0, \infty)^k, \prod_{j=1}^k x_j^{2\beta_j} dx)$.

On the other hand, we can write

$$H_{\beta_1, \dots, \beta_k}(g)(x) \leq \frac{1}{x_1^{2\beta_1+1}} \int_0^{x_1} \frac{1}{x_2^{2\beta_2+1}} \int_0^{x_2} \dots \frac{1}{x_k^{2\beta_k+1}} \int_0^{x_k} |g(y)| \prod_{j=1}^k y_j^{2\beta_j} dy, \quad x \in (0, \infty)^k.$$

Since, as it is well known, the Hardy type operator H_β given by, for every $\beta > -1/2$,

$$H_\beta(g)(x) = \frac{1}{x^{2\beta+1}} \int_0^x g(y) y^{2\beta} dy, \quad x \in (0, \infty),$$

is bounded from $L^p((0, \infty), x^{2\beta} dx)$ into itself, for every $1 < p < \infty$ (see [13]), from [4, Proposition 1] we deduce that $H_{\beta_1, \dots, \beta_k}$ is a bounded operator from $L^p((0, \infty)^k, \prod_{j=1}^k x_j^{2\beta_j} dx)$ into itself, for every $1 < p < \infty$.

Then, from (54) we infer that the operator $\mathcal{H}_{\beta_1, \dots, \beta_k}$ is bounded from $L^p((0, \infty)^k, \prod_{j=1}^k x_j^{2\beta_j} dx)$ into itself, for every $1 < p < \infty$, and from $L^1((0, \infty)^k, \prod_{j=1}^k x_j^{2\beta_j} dx)$ into $L^{1,\infty}((0, \infty)^k, \prod_{j=1}^k x_j^{2\beta_j} dx)$. Also, $S_{\lambda_1, \dots, \lambda_n}^{n, n, i}$ is bounded from $L^p((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$ into itself, for every $1 < p < \infty$, and from $L^1((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$ into $L^{1,\infty}((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$, for every $i = 1, \dots, n$.

Assume that $1 \leq i \leq k < n$. By using (51) and (53) we have that

$$\begin{aligned}
|S_{\lambda_1, \dots, \lambda_n}^{k, k, i}(g)(x)| &\leq C \int_0^{\frac{x_1}{2}} \dots \int_0^{\frac{x_k}{2}} \int_{2x_{k+1}}^\infty \dots \int_{2x_n}^\infty \int_0^\infty \frac{e^{-\left(\sum_{j=1}^k x_j^2 + \sum_{j=k+1}^n y_j^2\right)/20t}}{\sum_{t=1}^n (\lambda_j + 1/2) + 1} dt |g(y)| \prod_{j=1}^n y_j^{2\lambda_j} dy \\
&\leq C \int_0^{\frac{x_1}{2}} \dots \int_0^{\frac{x_k}{2}} \int_0^\infty \frac{e^{-\sum_{j=1}^k x_j^2/20t}}{\sum_{t=1}^k (\lambda_j + 1/2) + 1} dt \int_{2x_{k+1}}^\infty \dots \int_{2x_n}^\infty \frac{|g(y)|}{\prod_{j=k+1}^n y_j^{2\lambda_j+1}} \prod_{j=1}^n y_j^{2\lambda_j} dy \\
&\leq C \frac{1}{\left(\sum_{j=1}^k x_j^2\right)^{\sum_{j=1}^k (\lambda_j + 1/2)}} \int_0^{\frac{x_1}{2}} \dots \int_0^{\frac{x_k}{2}} \left(\int_{2x_{k+1}}^\infty \dots \int_{2x_n}^\infty \frac{|g(y)|}{y_{k+1} \dots y_n} dy_n \dots dy_{k+1} \right) \\
&\quad \times \prod_{j=1}^k y_j^{2\lambda_j} dy_k \dots dy_1, \quad x \in (0, \infty)^n.
\end{aligned}$$

According to [4, Proposition 1] and by taken into account the L^p -boundedness properties of the operator $H_{\lambda_1, \dots, \lambda_k}$ and $\mathcal{S}_{n-k}$ we conclude that the operator $S_{\lambda_1, \dots, \lambda_n}^{k, k, i}$ is bounded from $L^p((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$ into itself, for every $1 < p < \infty$, and from $L^1((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$ into $L^{1, \infty}((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$.

In a similar way we can see that if $0 < k < i \leq n$ the operator $S_{\lambda_1, \dots, \lambda_n}^{k, k, i}$ is bounded from $L^p((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$ into itself, for every $1 < p < \infty$, and from $L^1((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$ into $L^{1, \infty}((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$.

Let $1 \leq i \leq k < n$. By using (4), (5), (51) and (53), we get

$$\begin{aligned}
|S_{\lambda_1, \dots, \lambda_n}^{k, n, i}(g)(x)| &\leq C \int_0^{\frac{x_1}{2}} \dots \int_0^{\frac{x_k}{2}} \int_{\frac{x_{k+1}}{2}}^{2x_{k+1}} \dots \int_{\frac{x_n}{2}}^{2x_n} \int_0^\infty \frac{e^{-\sum_{j=1}^k x_j^2/20t}}{\sum_{t=1}^k (\lambda_j + 1/2) + 1} \\
(55) \quad &\times \prod_{j=k+1}^n \left(\frac{1}{x_j^{2\lambda_j+1}} + (x_j y_j)^{-\lambda_j} \frac{e^{-(x_j - y_j)^2/4t}}{\sqrt{t}} \right) dt |g(y)| \prod_{j=1}^n y_j^{2\lambda_j} dy, \quad x \in (0, \infty)^n.
\end{aligned}$$

Suppose that $k < l \leq n$, $l \in \mathbb{N}$, and define

$$\begin{aligned}
\mathbb{S}_{\lambda_1, \dots, \lambda_n}^{k, l}(g)(x) &= \int_0^{\frac{x_1}{2}} \dots \int_0^{\frac{x_k}{2}} \int_{\frac{x_{k+1}}{2}}^{2x_{k+1}} \dots \int_{\frac{x_n}{2}}^{2x_n} \int_0^\infty \frac{e^{-\left(\sum_{j=1}^k x_j^2 + \sum_{j=k+1}^l (x_j - y_j)^2\right)/20t}}{\sum_{t=1}^k (\lambda_j + 1/2) + 1 + (l-k)/2} dt \\
&\quad \times \prod_{j=k+1}^l (x_j y_j)^{-\lambda_j} \prod_{j=l+1}^n \frac{1}{x_j^{2\lambda_j+1}} g(y) \prod_{j=1}^n y_j^{2\lambda_j} dy, \quad x \in (0, \infty)^n.
\end{aligned}$$

We have that, for each $x \in (0, \infty)^n$,

$$\begin{aligned}
 |\mathbb{S}_{\lambda_1, \dots, \lambda_n}^{k, l}(g)(x)| &\leq C \int_0^{\frac{x_1}{2}} \cdots \int_0^{\frac{x_k}{2}} \int_{\frac{x_{k+1}}{2}}^{2x_{k+1}} \cdots \int_{\frac{x_l}{2}}^{2x_l} \frac{\prod_{j=k+1}^l (x_j y_j)^{-\lambda_j}}{(\sum_{j=1}^k x_j^2 + \sum_{j=k+1}^l (x_j - y_j)^2)^{\sum_{j=1}^k (\lambda_j + 1/2) + (l-k)/2}} \\
 (56) \quad &\times \left(\frac{1}{\prod_{j=l+1}^n x_j^{2\lambda_j + 1}} \int_{\frac{x_{l+1}}{2}}^{2x_{l+1}} \cdots \int_{\frac{x_n}{2}}^{2x_n} |g(y)| \prod_{j=l+1}^n y_j^{2\lambda_j} dy_n \cdots dy_{l+1} \right) \prod_{j=1}^l y_j^{2\lambda_j} dy_l \cdots dy_1.
 \end{aligned}$$

Assume that $r, s \in \mathbb{N}$, $0 < s < r$, and $\beta_j > -1/2$, $j = 1, \dots, r$. We consider the operator

$$\begin{aligned}
 Y_{\beta_1, \dots, \beta_r}^s(g)(x) &= \int_0^{\frac{x_1}{2}} \cdots \int_0^{\frac{x_s}{2}} \int_{\frac{x_{s+1}}{2}}^{2x_{s+1}} \cdots \int_{\frac{x_r}{2}}^{2x_r} \frac{\prod_{j=s+1}^r (x_j y_j)^{-\beta_j}}{(\sum_{j=1}^s x_j^2 + \sum_{j=s+1}^r (x_j - y_j)^2)^{\sum_{j=1}^s (\lambda_j + 1/2) + (r-s)/2}} \\
 &\times g(y) \prod_{j=1}^r y_j^{2\lambda_j} dy, \quad x \in (0, \infty)^r.
 \end{aligned}$$

By proceeding as in the proof of Case 3 in [15] we can see that the operator $Y_{\beta_1, \dots, \beta_r}^s$ is bounded from $L^1((0, \infty)^r, \prod_{j=1}^r x_j^{2\beta_j} dx)$ into $L^{1, \infty}((0, \infty)^r, \prod_{j=1}^r x_j^{2\beta_j} dx)$. Since the operator $Z_{\lambda_{l+1}, \dots, \lambda_n}$ is bounded from $L^1((0, \infty)^r, \prod_{j=1}^r x_j^{2\beta_j} dx)$ into itself, by [4, Proposition 1], we deduce from (56) that the operator $\mathbb{S}_{\lambda_1, \dots, \lambda_n}^{k, l}$ is bounded from $L^1((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$ into $L^{1, \infty}((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$.

On the other hand we also have that

$$\begin{aligned}
 |\mathbb{S}_{\lambda_1, \dots, \lambda_n}^{k, l}(g)(x)| &\leq C \int_0^{\frac{x_1}{2}} \cdots \int_0^{\frac{x_k}{2}} \int_0^\infty \frac{e^{-\sum_{j=1}^k x_j^2 / 20t}}{t^{\sum_{j=1}^k (\lambda_j + 1/2) + 1}} dt \left\{ \frac{1}{\prod_{j=l+1}^n x_j^{2\lambda_j + 1}} \int_{\frac{x_{l+1}}{2}}^{2x_{l+1}} \cdots \int_{\frac{x_n}{2}}^{2x_n} \right. \\
 &\left(\sup_{t>0} \int_{\frac{x_{k+1}}{2}}^{2x_{k+1}} \cdots \int_{\frac{x_l}{2}}^{2x_l} \prod_{j=k+1}^l (x_j y_j)^{-\lambda_j} \frac{e^{-(x_j - y_j)^2 / 4t}}{\sqrt{t}} |g(y)| \prod_{j=k+1}^l y_j^{2\lambda_j} dy_l \cdots dy_{k+1} \right) \\
 &\times \prod_{j=l+1}^n y_j^{2\lambda_j} dy_n \cdots dy_{l+1} \left. \right\} \prod_{j=1}^k y_j^{2\lambda_j} dy_1 \cdots dy_k \\
 &\leq C \frac{1}{\prod_{j=1}^k x_j^{2\lambda_j + 1}} \int_0^{\frac{x_1}{2}} \cdots \int_0^{\frac{x_k}{2}} \left\{ \frac{1}{\prod_{j=l+1}^n x_j^{2\lambda_j + 1}} \int_{\frac{x_{l+1}}{2}}^{2x_{l+1}} \cdots \int_{\frac{x_n}{2}}^{2x_n} \left(\sup_{t>0} \int_{\frac{x_{k+1}}{2}}^{2x_{k+1}} \cdots \int_{\frac{x_l}{2}}^{2x_l} \right. \right. \\
 &\times \prod_{j=k+1}^n (x_j y_j)^{-\lambda_j} \frac{e^{-(x_j - y_j)^2 / 4t}}{\sqrt{t}} |g(y)| \prod_{j=k+1}^l y_j^{2\lambda_j} dy_l \cdots dy_{k+1} \left. \right) \prod_{j=l+1}^n y_j^{2\lambda_j} dy_n \cdots dy_{l+1} \left. \right\} \\
 &\times \prod_{j=1}^k y_j^{2\lambda_j} dy_1 \cdots dy_k, \quad x \in (0, \infty)^n.
 \end{aligned}$$

According to the L^p -boundedness properties of the operators $\Omega_{\beta_1, \dots, \beta_r}$, $Z_{\beta_1, \dots, \beta_r}$, and $H_{\beta_1, \dots, \beta_r}$, $\beta_j > -1/2$, $j = 1, \dots, r$, and $r \in \mathbb{N}$, by using [4, Proposition 1], we obtain that $\mathbb{S}_{\lambda_1, \dots, \lambda_n}^{k, l}$ is

bounded from $L^p((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$ into itself, for every $1 < p < \infty$.

We conclude that $S_{\lambda_1, \dots, \lambda_n}^{k, n, i}$ is bounded from $L^p((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$ into itself, for every $1 < p < \infty$, and from $L^1((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$ into $L^{1, \infty}((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$, provided that $1 \leq i \leq k < n$.

From (37) and (44) we get for $\alpha > -1/2$

$$(57) \quad \left| \frac{\partial}{\partial t} W_t^\alpha(u, v) \right| \leq C \left((uv)^{-\alpha} \frac{e^{-(u-v)^2/10t}}{t^{3/2}} + \frac{e^{-(u^2+v^2)/10t}}{t^{\alpha+3/2}} \right), \quad t, u, v \in (0, \infty).$$

Suppose now that $1 \leq k < i \leq n$. By using (4), (5), (51) and (57) we obtain

$$\begin{aligned} |S_{\lambda_1, \dots, \lambda_n}^{k, n, i}(g)(x)| &\leq C \int_0^{\frac{x_1}{2}} \cdots \int_0^{\frac{x_k}{2}} \int_{\frac{x_{k+1}}{2}}^{2x_{k+1}} \cdots \int_{\frac{x_n}{2}}^{2x_n} \int_0^\infty \frac{e^{-\sum_{j=1}^k x_j^2/20t}}{\sum_{j=1}^k (\lambda_j + 1/2)} \\ &\quad \times \prod_{j=k+1, j \neq i}^n \left(\frac{1}{x_j^{2\lambda_j+1}} + (x_j y_j)^{-\lambda_j} \frac{e^{(x_j - y_j)^2/10t}}{\sqrt{t}} \right) \left((x_i y_i)^{-\lambda_i} \frac{e^{-(x_i - y_i)^2/10t}}{t^{3/2}} \right. \\ &\quad \left. + \frac{e^{-(x_i^2 + y_i^2)/4t}}{t^{\lambda_i+3/2}} \right) dt |g(y)| \prod_{j=1}^n y_j^{2\lambda_j} dy \\ &\leq C \int_0^{\frac{x_1}{2}} \cdots \int_0^{\frac{x_k}{2}} \int_{\frac{x_{k+1}}{2}}^{2x_{k+1}} \cdots \int_{\frac{x_n}{2}}^{2x_n} \int_0^\infty \frac{e^{-\sum_{j=1}^k x_j^2/20t}}{\sum_{j=1}^k (\lambda_j + 1/2) + 1} \\ &\quad \times \prod_{j=k+1}^n \left(\frac{1}{x_j^{2\lambda_j+1}} + (x_j y_j)^{-\lambda_j} \frac{e^{(x_j - y_j)^2/10t}}{\sqrt{t}} \right) dt |g(y)| \prod_{j=1}^n y_j^{2\lambda_j} dy, \quad x \in (0, \infty)^n. \end{aligned}$$

Then, as above, $S_{\lambda_1, \dots, \lambda_n}^{k, n, i}$ is bounded from $L^p((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$ into itself, for every $1 < p < \infty$, and from $L^1((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$ into $L^{1, \infty}((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$, when $1 \leq k < i \leq n$.

By proceeding in a similar way we can see that the operator $S_{\lambda_1, \dots, \lambda_n}^{k, l, i}$ is bounded from $L^p((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$ into itself, for every $1 < p < \infty$, and from $L^1((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$ into $L^{1, \infty}((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$, when $0 \leq k < l < n$, and $i = 1, \dots, n$.

By (49) we conclude that $S_{\lambda_1, \dots, \lambda_n}^{k, l}$, $0 \leq k \leq l \leq n$, $k \neq 0$ or $l \neq n$, is bounded from $L^p((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$ into itself, for every $1 < p < \infty$, and from $L^1((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$ into $L^{1, \infty}((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$. Then, the operator $\mathcal{G}_{\lambda_1, \dots, \lambda_n}$ has the same L^p -boundedness properties.

From (40) it follows that the maximal operator $T_{\lambda_1, \dots, \lambda_n}^{m, *}$ is bounded from $L^p((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$ into itself, for every $1 < p < \infty$, and from $L^1((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$ into $L^{1, \infty}((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$.

Thus the proof of this proposition is completed. $\square$

From Proposition 2.2 we can deduce by using standard arguments the following result.

Proposition 2.3. *Let $\lambda_j > -1/2$, $j = 1, \dots, n$. For every $f \in L^p((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$, $1 \leq p < \infty$, the limit*

$$\lim_{\varepsilon \rightarrow 0^+} \int_{(0, \infty)^n, |x-y| > \varepsilon} f(y) K_{\lambda_1, \dots, \lambda_n}^\phi(x, y) \prod_{j=1}^n y_j^{2\lambda_j} dy,$$

exists, for almost all $x \in (0, \infty)^n$. Here $K_{\lambda_1, \dots, \lambda_n}^\phi(x, y)$ is defined as in Proposition 2.2. Moreover, the operator $\mathbb{T}_{\lambda_1, \dots, \lambda_n}^m$ defined by

$$\mathbb{T}_{\lambda_1, \dots, \lambda_n}^m(f)(x) = \lim_{\varepsilon \rightarrow 0^+} \int_{(0, \infty)^n, |x-y| > \varepsilon} f(y) K_{\lambda_1, \dots, \lambda_n}^\phi(x, y) \prod_{j=1}^n y_j^{2\lambda_j} dy, \text{ a.e. } x \in (0, \infty)^n,$$

is bounded from $L^p((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$ into itself, for every $1 < p < \infty$, and from $L^1((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$ into $L^{1,\infty}((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$.

Since $C_c^\infty((0, \infty)^n)$ is a dense subspace of $L^p((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$, $1 \leq p < \infty$, it follows that,

for every $f \in L^2((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$,

$$T_{\lambda_1, \dots, \lambda_n}^m(f)(x) = - \lim_{\varepsilon \rightarrow 0^+} \left(\int_{(0, \infty)^n, |x-y| > \varepsilon} f(y) K_{\lambda_1, \dots, \lambda_n}^\phi(x, y) \prod_{j=1}^n y_j^{2\lambda_j} dy + \alpha(\varepsilon) f(x) \right), \text{ a.e. } x \in (0, \infty)^n,$$

where α is a bounded function on $(0, \infty)$, and $T_{\lambda_1, \dots, \lambda_n}^m$ can be extended from $L^2((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$

to $L^p((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$ as a bounded operator from $L^p((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$ into itself, for every $1 < p < \infty$, and from $L^1((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$ into $L^{1,\infty}((0, \infty)^n, \prod_{j=1}^n x_j^{2\lambda_j} dx)$.

The proof of Theorem 1.1 is finished.

REFERENCES

- [1] J.J. Betancor, T. Martínez, and L. Rodríguez-Mesa, *Laplace transform type multipliers for Hankel transforms*, Canad. Math. Bull., 51 (2008), 487-496.
- [2] J.J. Betancor, and L. Rodríguez-Mesa, *Weighted inequalities for Hankel convolution operators*, Illinois J. Math., 44 (2) (2000), 230-245.
- [3] J.J. Betancor, and K. Stempak, *Relating multipliers and transplantation for Fourier-Bessel expansions and Hankel transform*, Tohoku Math. J., (2) 53 (2001), no. 1, 109-129.
- [4] U. Dinger, *Weak type (1,1) estimates of the maximal function for the Laguerre semigroup in finite dimensions*, Revista matemática Iberoamericana, 8(1) (1992), 93-118.
- [5] J. García-Cuerva, G. Manceri, P. Sjögren, and J.L. Torrea, *Spectral multipliers for the Ornstein-Uhlenbeck semigroup*, J. Anal. Math., 78 (1999), 281-305.
- [6] G. Garrigós, and A. Seeger, *Characterizations of hankel multipliers*, Math. Annalen, 342 (2008), 31-68.
- [7] G. Gasper, and W. Trebels, *Multiplier criteria of Hormander type for Fourier series and applications to Jacobi series and Hankel transforms*, Math. Annalen, 242 (3) (1979), 225-240.
- [8] J. Gosselin, and K. Stempak, *A weak type estimate for Fourier-Bessel multipliers*, Proc. Amer. Math. Soc., 106 (3) (1989), 655-662.
- [9] D.L. Guy, *Hankel transformations and weighted p-norms*, Trans. Amer. Math. Soc., 95 (1960), 137-189.
- [10] R. Kapelko, *A multiplier theorem for Hankel transforms*, Revista Mat. Compl., 11 (2) (1998), 281-288.
- [11] N.N. Lebedev, *Special functions and their applications*, Dover publications, New York, 1972.
- [12] S. Molina and S. Trione, *On the n-dimensional Hankel transforms of arbitrary order*, Integral Transforms Spec. Funct., 19 (2008), no. 5-6, 327-332.
- [13] B. Muckenhoupt, *Hardy's inequality with weights*, Studia Math., 44 (1972), 31-38.
- [14] B. Muckenhoupt, and E. Stein, *Classical expansions and their relation to conjugate harmonic functions*, Trans. Amer. Math. Soc., 118 (1965), 17-92.
- [15] A. Nowak, and P. Sjögren, *Weak type (1,1) estimates for maximal operators associated with various multi-dimensional systems of Laguerre functions*, Indiana Univ Math. J., 56 (2007), 417-436.

- [16] A. Nowak, and K. Stempak, *Weighted estimates for the Hankel transform transplantation operator*, Tohoku Math. J., 58 (2006), 277-301.
- [17] E.M. Stein, *Topics in harmonic analysis related to the Littlewood-Paley theory*, Ann. of Math. Studies, vol. 63, Princeton Univ. Press, Princeton, NJ, 1970.
- [18] E.M. Stein, *Harmonic analysis: real-variable methods, orthogonality, and oscillatory integrals*, Princeton University Press, Princeton, NJ, 1993.
- [19] K. Stempak, *The Littlewood-Paley theory for the Fourier-Bessel transform*, Preprint 45, Mathematical Institute, University of Wroclaw, 1985.
- [20] K. Stempak, and J.L. Torrea, *Poisson integrals and Riesz transform for Hermite function expansions with weights*, J. Funct. Anal., 202 (2003), 443-472.
- [21] K. Stempak, and W. Trebels, *Hankel multipliers and transplantation operators*, Studia Math., 125 (1) (1997), 51-66.
- [22] G.N. Watson, *A treatise on the theory of Bessel functions*, Cambridge University Press, Cambridge, 1966.

DEPARTAMENTO DE ANÁLISIS MATEMÁTICO, UNIVERSIDAD DE LA LAGUNA, CAMPUS DE ANCHIETA, AVDA. ASTROFÍSICO FRANCISCO SÁNCHEZ, s/n, 38271 LA LAGUNA (STA. CRUZ DE TENERIFE), SPAIN
E-mail address: jbetanco@ull.es

DEPARTAMENTO DE ANÁLISIS MATEMÁTICO, UNIVERSIDAD DE LA LAGUNA, CAMPUS DE ANCHIETA, AVDA. ASTROFÍSICO FRANCISCO SÁNCHEZ, s/n, 38271 LA LAGUNA (STA. CRUZ DE TENERIFE), SPAIN
E-mail address: alecas000@gmail.com

INSTITUTO DE CIENCIAS MATEMÁTICAS, CONSEJO SUPERIOR DE INVESTIGACIONES CIENTÍFICAS, SERRANO 121, 28006 MADRID, SPAIN
E-mail address: jezebel.curbelo@icmat.es